

KRES'YANINOV, A.G.

RAYKHBAUM, Ya.D.; KRES'YANINOV, A.G.

Electric erosion of metals in spark discharge. Izv. AN SSSR
Ser.fiz.18 no.2:258 Mr-Apr '54. (MLRA 7:11)
(Spectrum analysis) (Electric spark)

Kresz, I.

Investigation of the bone development of Korean Children. p. 71

ANTHROPOLOGIAI KOZLEMENYEK. Budapest, Hungary. Vol. 2, no. 3/4, 1958

Monthly List of East European Accessions, (EEAI) LC, Vol. 9, No 1, Jan. 1960
Uncl.

KRESZ, MARIA

Magyar paraszttvisélet, 1820-1867

Budapest, Hungary. Akademiai Kiado, 1956. 231 p.

Monthly List of East European Accessions (EEAI), LC. Vol. 8, No. 9, September 1959
Uncl.

KRESZ, MARIA

Ungarische Bauerntrachten, 1820-1867. (Übers. von Jeannette Hajdu) Berlin, Henscheiverlag Kunst and Gesellschaft, 1957. 163 p. 96 plates (part col., in portfolio) (Hungarian peasant costumes, 1820-1867. In German, illus., map, bibl., facsim., tables)

SO: Monthly Index of East European Accessions (EEAI) LC. Vol. 7, no. 4, April 1958

KRESZ, Otto

Experiment milling achievements in connection with the Italian varieties of wheat grown in Hungary. Pecsí musz szeml 6 no.1:20-25 Ja-Mr '61.

KRESZ, Otto

The Flour Mill of Pecs. Pecs1 musz szoml 5 no.3:24-26 J1-S '60.

GEBALA, A.; HANICKA, M.; KRET, B.

Investigations on the sex chromation in oral mucosa smears in
children. Folia biol 8 no.1/2:97-104 '60. (EEAI 10:4)

1. Pediatric Clinic, Medical Academy, Krakow; head: Prof. Dr. T.Giza
and Department of Histology, Medical Academy, Krakow; head: Prof.
Dr. J.Ackermann.

(CHILDREN)
(SEX (BIOLOGY))
(CHROMATION)
(MUCOUS MEMBRANE)
(MOUTH)

POLAND / High Molecular Chemistry.

Abs Jour : Ref Zhur - Khimiya No 5, 1959, No. 18046
Author : Lasocki, Z.; Kret, Z.
Inst : Not given
Title : Hydrolysis and Condensation of the Bi-functional Monomers
of Silicones. II. Partial Hydrolysis of Methylene-
dimethoxysilane.
Orig Pub : Roczn. chem. 1958, 32, No 3, 657-659
Abstract : Partial hydrolysis of methylethyldimethoxysilane with
water solution of methanol and in the presence of NaOH
catalyst was conducted. Properties of the obtained
products are described. For Part I see Ref Zhur - Khimiya
1958, 83995.

Card 1/1
1291 - B
1015 - C
1226 D, E, F, G
1055
1228 - H, I

E N D

I - 3

KOVES, Elemér; ROMWALTER, Alfred; KRETAI, József, marnok; KARPAFI, Gyula

Sándor Deniflee, iron metallurgical engineer. technical director of the Csepel Metal Works, 1890-1959; obituary. Koh lap 93 no.2:66-67 F '60.

1. Femkohászati Szakosztály elnöke (for Romwalter). 2. Csepeli Femmu (for Kretai). 3. Csepeli Femmu szalaghengeresze (for Karpafi).

MACSAY, Jozsef; ARHOS, Frigyes; KULTAI, Jozsef

Society name: Koz. lap 97 no. 3 518 N 64

1. Editor-in-Chief "Kobaszati Lapoz" (for Arhosi).

SOV/62-58-8-3/22

AUTHORS: Rubinshteyn, A. M., Akimov, V. M., Kretalova, L. D.

TITLE: The Properties and the Structure of $\text{NiO-Al}_2\text{O}_3$ -Catalysts
(Svoystva i struktura $\text{NiO-Al}_2\text{O}_3$ katalizatorov) Note 2:
The Radiographic Investigation of the Influence of the
Interaction of the Components and of the Conditions of
Thermal Treatment on the Phase Composition and Crystalline
Structure (Sobshcheniye 2. Rentgenograficheskoye izucheniye
vliyaniya vzaimosoyedinyeniya komponentov i usloviy termicheskoy
obrabotki na fazovyy sostav i kristallicheskuyu strukturu)

PERIODICAL: Izvestiya Akademii nauk SSSR, Otdeleniye khimicheskikh nauk,
1958, Nr 8, pp. 929-936 (USSR)

ABSTRACT: The investigation of the activity and effective selectivity
in the reaction of the decomposition of $i\text{-C}_3\text{H}_7\text{OH}$ on $\text{NiO-Al}_2\text{O}_3$ -
catalysts described in the previous paper (Ref 1) furnished
the proof of the interaction of the components in solid
 $\text{NiO-Al}_2\text{O}_3$ -catalysts. Therefore it was assumed that they ex-
hibit a phase structure. This assumption needed, however,

Card 1/4

SOV/62-58-8-3/22

The Properties and the Structure of $\text{NiO-Al}_2\text{O}_3$ -Catalysts. Note 2: The Radiographic Investigation of the Influence of the Interaction of the Components and of the Conditions of Thermal Treatment on the Phase Composition and Crystalline Structure

checking and proving by means of physical methods of investigation. Especially the detailed radiographic investigation of the $\text{NiO-Al}_2\text{O}_3$ -catalysts could remove the deviation of the results (preliminary work of the authors and investigations carried out by Milligan and Merten (Ref 2), and Milligan and Richardson (Ref 3)). The first important result obtained from this work was that the authors found that among the catalysts investigated no amorphous ones were detected. Thus, the data supplied by Milligan and Merten could not be disproved in any way. They also found that the X-ray structural measurements showed the crystalline structure of the commonly precipitated catalysts $(\text{pH}8)\text{NiO-Al}_2\text{O}_3$ containing from 0 to 100 molar % of NiO (in contrast to those catalysts described by Milligan and Merten (Ref 2)). It was shown that the conditions of production exert a greater influence on the structure of the catalysts than the quantitative correlation of the components. It was also

Card 2/4

S07/62-58-8-3/22

The Properties and the Structure of $\text{NiO-Al}_2\text{O}_3$ Catalysts. Note 2: The Radiographic Investigation of the Influence of the Interaction of the Components and of the Conditions of Thermal Treatment on the Phase Composition and Crystalline Structure

found that catalysts of less than 50 molar % of NiO are monophase and do not have a free NiO . By measuring the parameter of the crystal lattice and of the occurring modifications in the concentration of NiO it was found that these monophase catalysts consist of a spinel solution of NiAl_2O_4 (in excess $\gamma\text{-Al}_2\text{O}_3$). The catalytic properties of $\text{NiO-Al}_2\text{O}_3$ were compared to the data of the phase and structural analyses. It turned out that there exist optimum parameters of the spinel lattice within the range of from 7.90 to 7.95 Å (for the dehydration). There are 1 figure, 3 tables, and 13 references, 9 of which are Soviet.

ASSOCIATION: Institut organicheskoy khimii im. N. D. Zelinskogo Akademii nauk SSSR (Institute of Organic Chemistry imeni N. D. Zelinskii, AS USSR)

Card 3/4

SOV/62-58-8-3/22

The Properties and the Structure of $\text{NiO-Al}_2\text{O}_3$ -Catalysts. Note 2: The Radiographic Investigation of the Influence of the Interaction of the Components and of the Conditions of Thermal Treatment on the Phase Composition and Crystalline Structure

SUBMITTED: March 8, 1957

Card 4/4

KRETA1003, L.D.

Authors: Rubin, A. M., Kiselev, E. I., Petrov, V. M.,
Petrov, N. S., Kiselev, L. D.
Title: Polymorphism and Catalytic Properties of Al_2O_3
Chemical: From the Journal of the USSR Academy of Sciences
Abstract: Polymorphic modifications of Al_2O_3 and their catalytic
properties were studied. Preparation of γ -, α -,
 χ -, κ -, θ -, δ - Al_2O_3 modification is given. It
was shown that formation of different Al_2O_3 modifications
depends not only on the thermal conditions of dehydration,
but also on the structure of the starting aluminum
hydroxide. The following table gives a series of conditions
are given

Card 1/3

Association: N. D. Zelinskii Institute of Organic Chemistry Academy
of Sciences USSR (Institut organicheskoy khimii imeni
N. D. Zelinskogo Akademii nauk SSSR)

Submitted: May 5, 1958

Card 3/3

RUBINSHTEYN, A.M.; PRIBYTKOVA, N.A.; AKIMOV, V.M.; KRETALOVA, L.D.;
KLYACHKO-CURVICH, A.L.

Effect of alkali metal oxides on the activity, selectivity, and
phase composition of binary catalysts based on Al_2O_3 . Izv. AN SSSR.
Otd.khim.nauk no.9:1552-1558 S '61. (MIRA 14:9)

1. Institut organicheskoy khimii im. N.D.Zelinskogo AN SSSR.
(Alkali metal oxides) (Catalysts)

KRECHETOV, M.

SKGN-6 corn planter. Nauka i zhizn' 28 no.4:78-79 Ap '61.

(MIRA 14:5)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut mekhanizatsii
sel'skogo khozyaystva.

(Planters (Agricultural machinery))

RADOJICIC, Bozidar; KRETIĆ, Bozidar

Manifestations of agranulocytosis following PAS administration.
Contribution to the injurious effect of PAS on the hematopoietic
system. Srpski arh. celok. lek. 87 no.10:922-927 0 '59.

1. Interna klinika Vojno-medicinske akademije u Beogradu, Načelnik:
puk. prof. dr Milan Arsenijević.
(AGRANULOCYTOSIS etiol.)
(PARAAMINOSALICYLIC ACID eff. inj.)

KRSTIC, M.

②
Nonspecific osteomyelitis in childhood. B Kovacek and M
Krstic (Hum. med. Wkly., 1953, 103, 516-518).-- A review with
illustrative case reports and 7 references.
W. R. BULL

KRSTIC, Miroslav, dr.

Postoperative complications following gastrectomy for gastroduodenal ulcer. Med.-arh., Sarajevo 8 no.2:37-47 Mar-Apr 54.

1. Hirurška klinika Medicinskog fakulteta u Sarajevu, saf. prof. dr. B.Kovacevic.

(PEPTIC ULCER, surg.
gastrectomy, partial, postop. compl.)

~~KRETIC, Miroslav~~

Contribution to the problem of traumatic epilepsy; case reports.
Med. glas. 8 no.11-12:448-450 Nov-Dec 54.

1. Hirurska klinika Medicinskog fakulteta u Sarajevu (sef prof.
dr. B.Kovacevic)
(EPILEPSY
traum.)

KRETIC, MIROSLAV

MIROSLAV, Kretic, Dr.

Historical data on the development of thoracic surgery of the
respiratory tract. Med. arh., Sarajevo 9 no.2:79-86 March-Apr
55.

(RESPIRATORY TRACT, surg.
thoracic, hist. (Ser))

(THORAX, surg.
in resp. tract dis., hist. (Ser))

(HISTORY, MEDICAL,
thoracic surg. in resp. tract dis., development.
(Ser))

VIROSLAV Kretic, Doc., Dr.

Problems of the pancreatic cysts. Med. arh., Sarajevo 9 no.3:
67-73 May-June 55.

1. Hirurska Klinika Univerziteta u Sarajevu (Sef: Prof. Dr.
B. Kovacevic).

(PANCREAS, cysts

caused by abdom. trauma. (Ser))

(ABDOMEN, wds. & inj.

causing pancreatic cyst. (Ser))

(WOUNDS AND INJURIES,

abdomen, causing pancreatic cyst. (Ser))

(CYSTS,

pancreas caused by abdom. trauma. (Ser))

KRETIC, Miroslav, Doc., dr.

Injurious effects of antibiotic therapy in surgery. Med. arch.,
Sarajevo 9 no.4:55-61 July-Aug 55.

1. Iz Hirurske klinike Med. fakulteta u Sarajevu. (Sef prof.
dr. B. Kovacevic).

(ANTIBIOTICS, ther. use,
in surg., inj. eff. (Ser))

(SURGERY, OPERATIVE,
use of antibiotics, inj. eff. (Ser))

KRETIC, Miroslav, Doc., dr.

Operative risks in struma of Basedow's type. Med. arh.,
Sarajevo 9 no.6:43-47 Nov-Dec 55.

1. (z Hirurske klinike Med. fakulteta, Sef prof. dr.
Blagoje Kovacevic).

(GOITER,

(HYPERTHYROIDISM, surg.

indic for hyperthyroid goiter. (Ser))

KRETIC, Miroslav, Doc., dr.

Interpretation of cerebral commotion syndrome in craniocerebral injuries. Med. arh., Sarajevo 10 no.2:23-38 Mar-Apr 56.

1. Iz Hirurske klinike Medic. fakult. Sarajevu (Seč. prof. dr. Blagoje Kovacevic).

(CRANIUM wds. & inf.

craniocerebral caused by trauma & causing concussion, pathophysiol. & ther. (Ser))

(BRAIN, wds. & inj.

same)

(WOUNDS AND INJURIES,

same)

KRETIC, Miroslav, Doc., dr.

Diagnosis and treatment of medulla spinalis. Med. arh.,
Sarajevo 10 no.3:21-31 May-June 56.

1. Iz Hirurske klinike Medicinskog fakulteta, Sef: prof. dr.
Blagoje Kovacevic.
(SPINAL CORD, dis.
diag. & management (Ser))

KRETIC, M.

Causes of late dehiscence of anastomotic sutures in esophageal resection.
Acta chir. jugosl. 4 no.3:221-228 1957.

1. Hirurška klinika Medicinskog Fakulteta u Sarajevu (Ref: prof. dr.
Blagoje Kovacevic)

(ESOPHAGUS, surg.

resection for cancer with anastomosis to stomach, late
rupt. of suture (Ser))

(STOMACH, surg.

anastomosis to esophageal stum after resection for
esophageal cancer, late rupt. of suture (Ser))

(SUTURES,

causes of rupt. of anastomotic suture in resection of
esophagus for cancer with anastomosis to stomach (Ser))

KRETIC, Miroslav, Doc., dr.

~~Modern neurosurgical diagnosis and prognosis of intracranial tumors.~~
Modern neurosurgical diagnosis and prognosis of intracranial tumors. Med. arh., Sarajevo 11 no.1:1-40 Jan-Feb 57.

1. Hirurška klinika Medicinskog fakulteta u Sarajevu. Sef: prof. dr. Blagoje Kovacevic.

(BRAIN NEOPLASMS,
intracranial, diag. & progn. (Ser))

KRETIC, Miroslav, doc. d-r

Significance of intracranial hypotension in general practice.
Med. arh., Sarajevo 12 no.2:23-28 Mr-Apr '59.

1. Hirurška klinika Medicinskog fakulteta u Sarajevu, šef: prof.
d-r B. Kovacevic.
(CEREBROSPINAL FLUID)

KRETIC, Miroslav, doc. d-r

Survey of therapy and prognosis in cerebral abscess. Med. arh.,
Sarajevo 13 no.4:27-39 JI-Ag '59.

1. Hirurška klinika Medicinskog fakulteta u Sarajevu, šef: prof.
d-r Blagoje Kovacevic.
(BRAIN ABSCESS)

KRETIC, Miroslav, doc. d-r

Complications in sub-occipital punctures. Med. arh., Sarajevo
13 no.6:51-55 N-D '59.

1. Kirurska klinika Medicinskog fakulteta u Sarajevu, sef: prof.
d-r B. Kovacevic.

(PUNCTURE compl.)

(BRAIN surg.)

KRETIC, M.; BESAROVIC, Z.; KOSAK, O.

Our first results with the application of hibernotherapy in severe and acute cerebrocranial injuries. Acta chir.iugosl. 7(8) no.3: 185-200 '60.

1. Kirurska klinika Medicinskog fakulteta u Sarajevu (v.d.sefa prof. dr F.Lukac)

(BRAIN wds & inj)

(HIBERNATION ARTIFICIAL)

KRETIC, Miroslav, doc. d-r; BOGDANOV, Branislav, d-r

Pathological and clinical contribution to mesenterial cysts. Med.arh.,
Sarajevo 14 no.3:31-36 My-Je '60.

1. Hirurska klinika Medicinskog fakulteta u Sarajevu (Sef: prof.
d-r Blagoje Kovacevic)
(MESENTERY dis)
(CYSTS)

KRETIC, Miroslav, doc. d-r

Specificity of intracranial expansive processes. Med.arh., Sarajevo
14 no.7:33-37 Ja '61.

1. Hirurska klinika Medicinskog fakulteta u Sarajevu (Sef: prof. d-r
Blagoje Kovacevic)
(INTRACRANIAL PRESSURE)

KRETIC, M.; BOGDANOV, B.; RIMSKI, B.

A case of *Cysticercus cellulose* in the central nervous system.
Med. arh. 15 no.4:1-9 J1-Ag '61.

1. Hirurska klinika Med. fakulteta u Sarajevu (v. d. sefa: Prof.
Feodor Lukac) Institut za patolosku anatomiju (v.d. sefa: Doc. dr
A.Nikulic).

(CYSTICERCOSIS case reports) (BRAIN dis)

1. Smallpox, Vol. 1, No. 3, Part 61, pp. 4-7.
2. Smallpox, Vol. 1, No. 3, Part 61, pp. 4-7.
3. Smallpox, Vol. 1, No. 3, Part 61, pp. 4-7.
4. Smallpox, Vol. 1, No. 3, Part 61, pp. 4-7.
5. Smallpox, Vol. 1, No. 3, Part 61, pp. 4-7.
6. Smallpox, Vol. 1, No. 3, Part 61, pp. 4-7.
7. Smallpox, Vol. 1, No. 3, Part 61, pp. 4-7.
8. Smallpox, Vol. 1, No. 3, Part 61, pp. 4-7.
9. Smallpox, Vol. 1, No. 3, Part 61, pp. 4-7.
10. Smallpox, Vol. 1, No. 3, Part 61, pp. 4-7.

RECEIVED

KRETIC, Vjerocka (Zagreb, Gunduliceva 2)

Labor productivity, an important indicator in the economic and social development of our country. Tesla 9 no.4:2-7 '62.

GENTN, S.A., KRETIMIN, A.A., KAZIMIRSEIY, Ya.M., spets.red.; VASII'YEVA, G.N.,
red.; YARGOV, E.M., tekhn.red.

[Practices of the Detchino factory in producing dehydrated potatoes]
Opyt Detchinskogo zavoda po proizvoizstvu sushennogo kartofelia.
Moskva, Pishchepromizdat, 1957. 17 p. (MIRA 11:5)
(Potatoes--Drying)

KRETININ, A.A.

New book on vegetable drying technology ("Industrial drying of vegetables and potatoes" by S.A. Genin, V.I. Knizhnik. Reviewed by A.A. Kretinin). Kuzn. ov.prou. 12 no.5:42-43 My '57.

(MLRA 10:8)

(Vegetables--Drying)

(Genin, S.A.) (Knizhnik, V.I.)

RUDZITSKIY, A.A.; RYBIN, N.S.; KRETININ, A.A.; CHERNOMORSKIY, G.A.,
spetsred.

[Automatic control of the process of drying on conveyer driers]
Avtomatizatsiia protsessa sushki na konveiernykh sushilkakh.
Moskva, Gos.nauchno-issl.in-t nauchn. i tekhn.informatsii, 1959.
9 p. (MIRA 13:6)

(Drying apparatus)

KRETININ, A.A.

Conveyer dryer and the design of its inclined belt. Kons. i ov. prom.
14 no. 2:21-22 F '59. (MIRA 12:3)

1. Detchinskiy molochno-ovoshchenushil'nyy zavod.
(Drying apparatus)

38169. KRETININ. G. A.

Nash opyt stroitel'stva purdov i vodoyemov (Lesozashchitnaya
stantsiya im. Gor'kogo. Tamb. obl.) Mekhanizatsiya trudoyenkikh
i tyazhelykh rabot, 1949, no. 12, s. 31

KRETININ, N. I.

Our experiment in cross-cultivating sugar beets. Moskva, Gos. izd-vo sel'khoz. lit-ry, 1954. 8 p. (Biblioteka obmena opytom peredovikov sel'skogo khoziaistva) (54-43476)

SF220.R9K7

KRETININ, S. A.

USSR/Chemical Technology. Chemical Products and Their
Application - Silicates. Glass. Ceramics. Binders.

I-9

Abs Jour : Referat Zhur - Khimiya, No 4, 1957, 12512

Author : Kretinin S.A.

Inst : Voronezh University

Title : Investigation of the Effect of Chemical Additives HCl,
H₂SO₄, NaOH and K₂CO₃ on SEDIMENTATION of Sand with Clay

Orig Pub : Tr. Voronezhsk. un-ta, 1955, 35, 59-74

Abstract : Using two specimens of sand and a specimen of hydronica-
ceous clay from Western Kazakhstan, a study was made of
the optimal conditions of sedimentation (rate of inflow
and turbidity of slurry), and investigations were con-
ducted of changes in filtration coefficient and thick-
ness of the silt film on introduction into the same slur-
ry of K₂CO₃, HCl, H₂SO₄, NaOH and methylene blue (over
the range of 0.001 -0.1 N). Most effective are additions
of methylene blue (0.0003%), 0.01 N and 0.001 N K₂CO₃ and
0.005 N H₂SO₄.

Card 1/1

- 64 -

KRETININ, S.A.

Study of filtration during the mutual coagulation of sols.
Trudy VGU 57:169-175 '59. (MIRA 13:5)
(Colloids) (Filters and filtration)

L 141-64
ACCESSION NR: AR3006942 EPR/ENP(j)/EPF(c)/ENT(m)/BDS AFFTC/ASD Pe-4/PC-4/Pr-4 RM/WW/
S/0081/83/000/010/0698/0598 MAY

SOURCE: Rzh. Khimiya, Abs. 107499

AUTHOR: Mikhant'yev, B. I., Kretinin, S. A., Shatalov, V. P.

TITLE: Study of the properties of divinyl-styrene rubbers filled in the latex stage

CITED SOURCE: Tr. Labor. khimii vyssokomolekul. soyedeneniy. Voronezhsk. un-t, vy*p. 1, 1962, 162-169

TOPIC TAGS: Divinyl-styrene rubber, latex stage, rubber

TRANSLATION: A study was made of the conditions of filling SKS-3CAR with HAF carbon black, channel carbon black, Al sub 2 O sub 3, PN-6 oil, auto scrap-18, and mazut at the latex stage and on rollers. Carbon black dispersions were prepared with a magnetic striction vibrator with a frequency of 25 kilocycles (concentration of carbon black of 15%, vibration time of 20 minutes). With

Card 1/2

L 141-64

ACCESSION NR: AR3006942

the introduction of 0.2-0.5% leucanol the vibration time is lowered to 5-10 minutes. The combination of latex with the dispersion of carbon black and the oil emulsion was also conducted through vibration for 3-5 minutes. The mixture was coagulated by CaCl sub 2 with H sub 2 SO sub 4 or CH sub 3 COOH. The expenditures per ton of commercial rubber with HAF carbon black were: CaCl sub 2 -- 30 kg, CH sub 3 COOH -- 1.9 kg; with channel carbon black; CaCl sub 2 -- 15.6 kg, CH sub 3 COOH -- 8 kg. The product which was obtained was dried at 80-90 degrees with forced ventilation. Upon introducing the carbon black into the latex a more plastic mixture was obtained which yielded stronger and more elastic vulcanized rubbers; the speed of vulcanization was increased. Dispersions with leucanol yielded better rubbers than without it. A basic technological plan for the production of carbon black-butyric rubbers was proposed. A 20% aqueous solution of Al sub 2 O sub 3 was prepared in a ball mill (30 rev/min) for 3 hours at about 20 degrees. The expenditure of CaCl sub 2 for the coagulation of 1 ton of commercial rubber was 47 kg. There was no loss of Al sub 2 O sub 3 during the coagulation of the latex mixture. The introduction of Al sub 2 O sub 3 into the latex produced more plastic mixtures and stronger vulcanized rubbers than when it was introduced on rollers. G. Chasovshchikov

DATE ACQ: 01Jul63

SUB CODE: CH, MA

ENCL: CO

Card 2/2

KUZNETSOV, Viktor Vasil'yevich; KASATOCHKIN, V.I., retsenzent;
KRETININ, S.A., retsenzent; PALKINA, N.A., retsenzent;
KONDRASHKOVA, S.F., red.

[Physical and colloid chemistry] Fizicheskaya i kolloid-
naya khimiya. Moskva, Vysshaya shkola, 1964. 385 p.
(MIRA 17:5)

L 37023-65 EWT(m)/EPF(a)/ENP(j)/T Po-4/Pr-4 RM
 ACCESSION NR: AR5003002 S/0081/64/000/019/H083/H083

SOURCE: Ref. zh. Khimiya, Abs. 19zh274

AUTHOR: Mikhant'yev, B. I.; Kretinin, S. A.; Shirokov, Yu. P.

TITLE: Synthesis and polymerization of some organotitanium compounds with unsaturated substituents

CITED SOURCE: Tr. Labor. khimii vysokomolekul. soyedineniy. Veronezhsk. un-t, vyp. 2, 1963, 47-49

TOPIC TAGS: heteroorganic compound, organotitanium compound, titanium polymer, butadiene polymer, isoprene polymer, titanium polyolefin, titanium tetrachloride, alkylene orthotitanate, orthophosphate initiator, thermal polymerization

TRANSLATION: The α -oxide of butadiene, boiling point 66-68C, n_{D}^{20} 1.4160, and the α -oxide of isoprene, boiling point 78-80C, n_{D}^{20} 1.4140, were obtained by the usual methods. By adding CH_3OH to these compounds in the presence of $BF_3 \cdot O(C_2H_5)_2$, the authors then obtained 2-methoxybuten-3-ol-1, boiling point 141-143C, n_{D}^{20} 1.4290, and 2-methoxy-2-methylbuten-3-ol-1, boiling point 75-77C/50 mm, n_{D}^{20} 1.4385. In an atmosphere of N_2 , 3.5 g of $TiCl_4$, boiling point 136C, were gradually

Cont 1/2

L 37023-65
ACCESSION NR: AR5003002

added to 150 ml of absolute benzene at 0C, the mixture was saturated with dry NH_3 , and the suspension of $\text{TiCl}_4 \cdot 8\text{NH}_3$ which was formed was gradually treated with a 1.5-fold excess of 2-methoxybuten-3-ol-1; after 5 hours at $\leq 10^\circ\text{C}$, the filtrate was evaporated off at 450 mm and (2-methoxybuten-3)-orthotitanate, $\text{C}_{20}\text{H}_{36}\text{O}_8\text{Ti}$ (I), boiling point 162-163C/2-2.5 mm, n_D^{20} 1.5050, d_4^{20} 1.0895, was obtained from the residue in 52% yield. Analogously, (2-methoxy-2-methylbuten-3)-orthotitanate, $\text{C}_{21}\text{H}_{38}\text{O}_8\text{Ti}$ (II), boiling point 152-154C/2-2.5 mm, n_D^{20} 1.5260, d_4^{20} 1.1190, was obtained from 2-methoxy-2-methylbuten-3-ol-1 in 35% yield. At 20-25C with orthophosphoric acid as an initiator, the authors obtained a polymer from I having a melting point of 240-250C and a polymer from II having a melting point of 250-255C. The thermal polymerization of I and II (at 150-180C for 2 hours) yielded brittle films with low adhesion to glass and softening points of 375 and 360-380C, respectively. The thermal polymerization of I and II at 100C for 8 hours was unsuccessful; FeCl_3 , BF_3Na and $(\text{C}_6\text{H}_5\text{CO})_2\text{O}_2$ were found to be ineffective as initiators. F. Velichko.

SUB CODE: OC

ENCL: 00

Card 2/2 *lo*

L 37019-65 EMT(m)/EPP(c)/EPR/EMP(j) Pc-4/Pr-4/Pe-4 NW/RM
 ACCESSION NR: AR5003012 S/0081/64/000/020/S082/S082

SOURCE: Ref. zh. Khimiya, Abs. 208511

AUTHOR: Mikhant'yev, B. I.; Kretinin, S. A.; Gostev, M. M.; Shatalov, V. P.;
 Markina, E. I.; Senyuk, Ye. P.

TITLE: Butadiene-styrene rubbers filled with carbon black and oil and produced by high-temperature polymerization

CITED SOURCE: Tr. Labor. khimii vysokomolekul. soedineniy. Voronezhsk. un-t, vyp. 2, 1963, 103-108

TOPIC TAGS: synthetic rubber, butadiene rubber, styrene rubber, carbon black filler, gas black filler, channel black filler, oil filled rubber, high temperature polymerization, rubber mechanical property, rubber emulsifier, synthetic fatty acid, colophony, latex coagulation

TRANSLATION: The authors studied the properties of butadiene-styrene rubbers of the SKS-30 type, produced by high-temperature polymerization with the addition of 17.6-50.0 parts by weight PN-6181 and 50.0 parts by weight gas black, channel black or NAF black to latex stage. The following combinations were tested as

Card 1/2

L 37019-65

ACCESSION NR: AR5003012

emulsifying agents: Nekal and the Na soaps of synthetic fatty acids; Nekal and the K soaps of synthetic fatty acids; the K soap of hydrogenated colophony and the K soaps of synthetic fatty acids. The 20% carbon black dispersions were prepared by grinding in a ball mill for 24 hrs. in the presence of 4-6 parts by weight leukanol and 0.6 parts by weight NaOH (in relation to the carbon black). The oil emulsion was of commercial origin. During the coagulation of mixtures from Nekal latex, the best results were produced by CaCl_2 and CH_3COOH ; in the case of latex produced with the soaps of synthetic fatty acids, the best results were produced by a mixture of CaCl_2 , NaCl and CH_3COOH ; in the case of colophony latex, NaCl and H_2SO_4 gave the best results. During deformation of the initial rubber with 4500 g. raw mixtures of rubber filled with carbon black and oil (SMK rubber) had a somewhat greater plasticity and less reducibility than when carbon black was added to oil-filled rubber on the rollers. The strength of the SMK vulcanates was somewhat lower, however. The method of introducing the carbon black had no significant effect on the properties of rubber mixtures and vulcanates in soft rubber. The properties of rubber do depend, however, on the method of coagulation. The instantaneous (single-stage) coagulation of SMK rubber resulted in somewhat more rigid mixtures with increased strength and decreased relative elongation. A. Shvarts.

SUB CODE: MT
Card 2/2

ENCL: 00

L 22029-66 EWT(m)/EWP(j)/T GS/RM

ACC NR: AT6005937

(A)

SOURCE CODE: UR/0000/63/000/000/0047/0049

AUTHORS: Mikhant'yev, B. I. ; Kretinin, S. A. ; Shirokov, Yu. P.



ORG: Laboratory for the Chemistry of High-Molecular-Weight Compounds, Voronezh State University (Laboratoriya khimii vysokomolekulyarnykh soyedineniy Voronezhskogo gosudarstvennogo universiteta)

TITLE: Synthesis and polymerization of certain titanooorganic compounds containing unsaturated radicals

SOURCE: Voronezh. Universitet. Laboratoriya khimii vysokomolekulyarnykh soyedineniy. Trudy, no. 2, 1963. Monomery, khimiya i tekhnologiya SK (Monomers, chemistry, and technology of synthetic rubber), 47-49

TOPIC TAGS: organometallic compound, organotitanium compound, titanium compound, organic synthetic process, titanium, polymerization

ABSTRACT: The organotitanic compounds



Card 1/2

L 22029-66

ACC NR: AT6005937

0
were synthesized to extend the work of S. V. Nogina, R. Kh. Froydlina, and A. N. Nesmeyanov (Izv. AN SSSR, OKhN, 3, 327, 1950). The compounds were synthesized by reacting $TiCl_4 \cdot 8NH_3$ in dry benzene with 2-methoxybuten-3-ol-1 and with 2-methoxy-2-methylbuten-3-ol-1 respectively. The intermediate products were synthesized after the method of A. A. Petrov (ZhOKh, 11, 991, 1941; ZhOKh, 16, 1625, 1946) and of A. N. Pudovnik and S. G. Denislamova (ZhOKh, 27, 2363, 1957). Reaction yields and the characteristic physical constants for the synthesized compounds are tabulated. The polymerization of the synthesized compounds was studied. Only thermopolymerization and polymerization induced by orthophosphoric acid yielded polymers. Metallic sodium, BF_3 , $FeCl_3$, and benzoyl peroxide did not induce polymerization. The polymers obtained were ruby-red in color, brittle, and showed a poor adhesion to glass. Orig. art. has: 2 formulas.

SUB CODE: 07/

SUBM DATE: none/

ORIG REF: 007

OTH REF: 001

Card. 2/2dda

L 40962-66 ENT(m)/ENP(j) IJP(c) RM/JND

ACC NR: AR6016972 (A)

SOURCE CODE: UR/0081/65/000/024/S077/S078

AUTHOR: Mikhant'yev, B. I.; Gostev, M. M.; Kretinin, S. A.

TITLE: Carbon black-oil filled butadiene styrene rubber of low temperature polymerization obtained in a system with a Trilon Rongalite activating group

SOURCE: Ref. zh. Khimiya, Abs. 24S547

REF SOURCE: Tr. Labor. khimii vysokomolekul. soyedineniy. Voronezhsk. un-t, vyp. 3, 1964, 186-190

TOPIC TAGS: butadiene styrene rubber, filler, carbon black, polymerization catalyst, elasticity, tensile strength

ABSTRACT: The possibility of using channel and gas furnace blacks in reinforcement in SKS-30 ARK latex prepared with the Trilon Rongalite activating group was investigated. The carbon blacks were introduced into the latex as 20% dispersions stabilized with K-soaps of hydrated or disproportionated rosin. The following proportions of stabilizer were necessary to obtain stabilized dispersions: for channel black 4-5 parts by weight, for furnace black 3.5 parts by weight, for their mixtures (1:1) 5-6 parts by weight. Introduction of both carbon blacks and their

Card 1/2

L 40962-56

ACC NR: AR6016972

0
mixtures to the latex gives vulcanizates with lower modulus and greater elasticity. The strength of vulcanizates with furnace black is higher and with channel black it is lower than when the carbon black is added on the rolls. M. Ayzinson. [Translation of abstract].

SUB CODE: 07, 11, 10

Card 2/2 mlp

STEPCHKOV, K.A.; KRETININA, L.V.; ADAMSON, R.P., otv. za vyp.;
BERENSHTEYN, R.Ye., otv. za vyp.; MAHVELOVA, Ye.S.,
tekhn. red.

[Production of potato granules] Proizvodstvo kartofel'noi
krupki. Moskva, TSintipishcheprom, 1963. 24 p.
(MIRA 17:1)

(Potatoes, Drying)

KRETININA, T.I., vrach

Significance of formal gelification of the blood serum in liver diseases. Zdrav. Kazakh. 18 no. 2:49-52 '58. (MIRA 13:8)

1. Iz Alma-Atinskoy gorodskoy ob'yedinennoy klinicheskoy bol'nitsy.
(LIVER--DISEASES) (SERUM)

KRETININA, Yu.S.; PERESLEGIN, Yu.A.

Semigraphical method of calculating long-distance power transmission
lines with consideration of the loading effect. Obshch. energ.
no.3:34-38 '60. (MIRA 14:3)
(Electric power distribution)

Kretinsky, B.

A mixer for foamed concrete p. 156

(Stavivo. Vol. 35, no. 3, Mar. 1957. Praha, Czechoslovakia)

SO: Monthly List of East European Accessions (EEAL) LC, Vol. 6, no. 10, October 1957. Uncl.

CH

7

V. Methods of investigating phosgene. A. E. Kargin. *Zhur. Prikladnoi Khimii* 2, 483, 601 (1929). -- Less than 0.1% CO or CO₂ is sol. in COCl₂ up to -10°. Solv. of Cl₂ is 3.0-4.3% at -10° and of HCl -1.6-2.2% at 0° and 4% at -12°. COCl₂ does not corrode pure Fe even in the presence of Cl₂ but impure Fe rapidly forms FeCl₂.

Com. COCl₂ contains 0.08-0.20% FeCl₂, 0.2-1.2% HCl and 0.1-0.3% org. compds. But sometimes considerably more. For detection of COCl₂, aniline water (not very sensitive), filter paper wetted in a soln. of 1 g. *p*-dimethylaminobenzaldehyde and 1 g. diphenylamine in 20 g. EtOH (reacts with other compds. besides COCl₂) or in a soln. of 0.3 g. 1,3,6-nitroindolethylaminophenol and 0.25 g. *p*-dimethylaminophenol in 100 cc. CCl₄ (sensitivity less than usually reported and is affected by purity of reagents) can be used. The sp. gr. is 1.428-1.431 (little affected by the presence of 2-3% impurities) and is detd. with a hydrometer at 0°. Non-volatile impurities are found by vaporizing COCl₂ on a water bath. Removal of org. impurities is essential for the detn. of the FeCl₂ present. Cl₂ imparts to COCl₂ a yellow-greenish tinge, FeCl₂ a grayish yellow tinge (due to formation of unstable compds.), and org. compds. darken its color. For quant. analysis samples are taken by cooling the container and capillary tubing leading from it with a mixt. of Et₂O and solid CO₂. From 10 to 20 cc. is drawn into a small flask with 2 openings: one for filling and the other, covered with a sheet of rubber which has a small opening in the middle, for drawing the sample into the ampule. This flask is kept at -10-15°, and the ampule of about 0.5 cc. capacity is half filled through the opening in the rubber by cooling its bulb in a mixt. of Et₂O and solid CO₂. The ampule is sealed while its bulb is kept at -10°. In absence of Cl₂ or large amts. of HCl cooling with solid CO₂ is not required. The method of Kling and Schmitz (*J.* 13, 1088, 2058) is recommended. Its accuracy, contrary to general belief, is not affected by the presence of up to 6.5% of Cl₂. COCl₂ can be also analyzed by sapon. 0.25-0.35 g. of it with 25 cc. N NaOH soln. The mixt. is kept for 1-2 hrs. at 0° in a glass stoppered flask, diluted to 100-250 cc. with water, and titrated in a closed flask with

OVER

agitation by means of CO₂-free air. Greater accuracy is attained, and the result is not affected by the impurities present in COCl₂, by titration at 0° and comparison of the resulting colors in a colorimeter with those of standard solns. Free Cl₂ is detd. by mixing COCl₂ with a 0.1% KI soln. and titrating the liberated I₂ with 0.05 N 0.01 N Na₂S₂O₃. Instead of KI a 0.5% soln. of Na or K arsenite in the presence of an excess of bicarbonate might be used. HCl is detd. by means of Hg(CNS)₂ or Hg(CN)₂. About 1 g. of powd. Hg(CN)₂ is placed in a 500-cc. Drexel flask, the ampule contg. COCl₂ (or HCl) in abs. EtOH soln. is then introduced, and the flask evacuated. The ampule is broken by shaking and the mixt. is allowed to stand for 12-16 hrs. The air dried with H₂SO₄ is then soaked through the flask for 8-10 hrs. and bubbled through a 2 N soln. of alkalis which is afterward titrated. This method is, however, not an exact one. Analysis of COCl₂ by detg. the amt. of Cl₂ formed on heating COCl₂ with powd. metals is inaccurate. Analysis by means of NaI and Me₂CO is good, but expensive, as very pure reagents should be used. HCl cannot be quantitatively sep'd from COCl₂ by washing with water.

V. KALICHURSKY

ca

Preparation of fatty, fatty-aromatic and fatty-naphthenic esters. A. KERNY AND
BAKAKINA. *Zhur. Prikladnoi Khim.* 2, 809-16(1929).--Prep. $(\text{MeO})\text{CO}(\text{OCH}_2\text{Ph})$
by heating, for $4\frac{1}{2}$ hrs. on a water bath and then for $\frac{1}{2}$ hr. on a gas burner, a mixt. of 200
g. benzyl alc., 90 g. pptd. chalk and 150 cc. MeO CO Cl . Remove CaCl_2 formed with
water and excess chalk with HCl . Purify by distg. under reduced pressure (b.p. 140-
141°) not to decomp. the product. The yield is 100%. Prep. $\text{Et CO}(\text{OC}_6\text{H}_5)_2$ (b.p.
148-154°) in a similar manner (80-85% yield) as well as $\text{iso Bu CO}(\text{OC}_6\text{H}_5)_2$ (b.p.
172-174°, d₄ 1.075; 87% yield). The same method is applicable to cyclohexanol
derivs. Prep. $(\text{MeO})\text{CO}(\text{OC}_6\text{H}_5)_2$ by heating, for 6 hrs. on a water bath, a mixt. of 50 g.
cyclohexanol, 40 cc. MeO CO Cl and 25 g. pptd. chalk. Wash with water, dry and
distil under vacuum. b.p. 190-200°. The yield is 50-55%. Use a similar procedure
for preps. $\text{Ri CO}(\text{OC}_6\text{H}_5)_2$ (b.p. 213.5-215.5°, 50% yield), $\text{iso Bu CO}(\text{OC}_6\text{H}_5)_2$ (b.p.
192°, b.p. 210-211°, 54.8% yield) and $\text{iso Am CO}(\text{OC}_6\text{H}_5)_2$ (b.p. 151°, b.p. 231.5-
251°; 62.4% yield).
V. KALICHEVSKY

ASB-5LA DETAILING LITERATURE CLASSIFICATION

Bis[phenylacetonitrile] sulfide, phenylacetonitrile mercaptan, phenylacetonitrile thiocyanate and their derivatives. A. KANTU and A. PANCHENKO. *J. Russ. Phys.-Chem. Soc.* 61, 1975-93 (1929).—The aliphatic-aromatic sulfides and thiocyanates have here an entirely new field. **Bis[phenylacetonitrile] sulfide (I)**, $\text{PhCH(CN)}_2\text{S}$, was obtained (1) by adding gradually while cooling 132 g. of NaS in 400 cc. of alc. to 150 g. of PhCH(CN)_2 (II) in 180 cc. of alc., then boiling, allowing to stand 2 days, filtering from CaH_2 to remove dicyanobisulfide (III) (yield about 20%), m. 132°. From the $\text{PhCH(CN)}_2\text{SH}$, I and III. $\text{PhCH(CN)}_2\text{Br}$ (V) under these conditions gives I greatly contaminated with III, IV and S. (2) To 50 g. of V in 40 cc. of alc. was added 40 g. of NaS and 63 g. of S in 200 cc. of alc., filtered the next day, the crystals washed in a Buchner with alc. and H_2O , added to I sepd. from the mother liquor, recrystd. twice from alc., m. 180°, yield 7.8 g. (22.3%), light yellow crystals, insol in H_2O and ligroin, sparingly sol. in alc., MeOH, AcOH, easily sol. in C_6H_6 and AcOH. The constitution is proved by the method of prepn. and by sapon., whereby first is formed III and then over, the reduction with HI gives the dicyanobisethyl (VIII), [PhCH(CN)]_2 , with the lower m. p., while the higher melting stereoisomer was [prepd. by reducing I with nascent H. (*Ber.* 25, 258, 294 (1902); 37, 4067 (1904); *J. Chem. Soc.* 83, 998, 1003 (1911)). I gives with NH_4OH no amine, but heating I in dil. alc. with NH_4OH HCl while passing CO_2 into the liquid produces a good yield of cranostillene-carbozamide, $\text{PhC(CN).C(CN)H}_2\text{Ph}$, which is converted with KOH to IV. **Prepn. of IV.** (1) To 39.2 g. of V in 200 cc. of alc. is added a warm soln. of 80 g. of $\text{Na}_2\text{S}_2\text{O}_3$, the mixt. is refluxed for 2.5 hrs., the greater part of the alc. expelled, cooled, decanted, with 150 cc. of 2 N HCl, the layer of oil sepd., dissolved in hot alc., filtered from S and allowed to stand for a few days, m. 101°, yield 11 g. (30.9%) (2) To 30 g. of II in 30 cc. of alc. was slowly

OVER

added while cooling 9 g of cryst. Na_2S in 120 g of alc. satd. with H_2S . Filtered the next day, the filtrate dild. with 2 vols. of H_2O , heavy yellow oil, yield 20.7 g. (91.35%). When working with 80 g. of V there is obtained 19.5 g. (51.0%) of crude IV and 11 g. (32.7%) of crude I. IV is a sirupy mass of crystals, m. 101° , faint odor, insol. in H_2O and petroleum ether, sol. in alc., Et_2O and C_6H_6 ; it shows double mol. wt., is decompd. on boiling with KOH to H_2S and VI, while reduction with III in AcOH produces VII. *Phenylacetonitrile thiosemide*. To 19.5 g. of V in 20 cc. of alc. is added 8.5 g. of NH_4CNS in 35 cc. of warm alc., the mixt. brought to a boil, filtered from NH_4Br , the filtrate cooled, dild. with H_2O , yield 14 g. (80.4%); recrystd. 4 times from a mixt. of 1 part of C_6H_6 and 2 parts of ligroin, white, odorless needles, m. 63.5° , sol. in C_6H_6 and alc., insol. in H_2O , petroleum ether, decompd. on distn. at 130° at atm. pressure and is *durio* to H_2S and resinous matter; on sapon. with alc. KOH it gives KCNS and III, which is decompd. with KOH into NH_3 and VI; reducing with III in AcOH produces a little of VII and resinous matter.

CHAS. H. BANC

10

Ca

The action of zinc dust and zinc oxide on the halogen derivatives of the aliphatic series. A. B. KAMRUV. J. Russ Phys Chem Soc. 61, 2143 (1920). The well known reactions between halogen substituted hydrocarbons and their deriva with Zn dust in presence of 98% or dl. alc result in (1) substitution of H for halogen with formation of hydrocarbons, (2) splitting off of 2 halogens with formation in the aliphatic series of cyclic compds., and in alicyclic series of unsatd. or bicyclic compds. (3) splitting off of H halide with formation of unsatd. compds. The mechanism of the reaction progresses in more simple cases with formation of intermediary zinc org compds. (Helfrich and Reid, C. A. 16, 2460; Kermack, et al., C. A. 16, 4201, Bales and Nickelson, C. A. 18, 51). The action of Zn dust on halogen deriva compd. S is here investigated. The interaction between $(C_2H_5CH_2)_2S$ (I), and Zn dust is a complex reaction resulting in a large no. of compds. The I is entirely decompd. with formation of C_2H_4 , H_2S , diethylene disulfide $S(C_2H_5CH_2)_2$ (II), HCl , etc. $2I + 2Zn = II + 2ZnCl_2 + 2C_2H_4$; $I + Zn = II + (CH_3CH_2)_2S$ (III), and $I + Zn + H_2 = H_2S + ZnCl_2 + 2C_2H_4$. The decompn. is caused by metallic Zn and not by ZnO present as impurity. Among liquid products of the reaction are found some R_2S , $CH_3CH_2SCH_2CH_2CH_3$ (IV) and $(CH_3CH_2)_2SH$. The formation of the first 2 is explained by addn. of H to III: $III + 2H_2 = R_2S$ and $III + H_2 = IV$. III and its polymers are formed in large amts. when I is heated with metals and their oxides in presence of inert solvents, such as xylene, and not of alc; a part of III is hereby converted to II, which is completely decompd. to gaseous prodn. Some tests produced $CH_3CH_2SCH_2CH_2CH_2OH$ III + $H_2O = CH_3CH_2SCH_2CH_2CH_2OH$. The main products of the reaction between I and Zn in the presence of alc are ethers. The main products of the general formula $(ROCH_2CH_2)_2S$, which are stable liquid compds. of thiodiglycol. under atm. pressure without decompn., the formation of which is thus capable of distin. under atm. pressure without decompn., the formation of which is thus formulated: $I + Zn = III + ZnCl_2 + H_2$; $I + ZnO = III + ZnCl_2 + H_2O$ and $III + 2ROH = (ROCH_2CH_2)_2S$. The other interpretation is based mainly on sapon. in presence of H_2O and alc: $(C_2H_5CH_2)_2S + H_2O + ZnCl_2 = C_2H_5CH_2OH + ZnCl_2$. $I + EtOH$

ASD-34 DETAIL LITERATURE CLASSIFICATION

OVER
CARD NO. 1

+ ZnO = $\text{HO}(\text{C}_2\text{H}_5\text{SC}_2\text{H}_4\text{OR}) + \text{ZnCl}_2$; and $\text{I} + 2\text{R}(\text{OH}) + \text{ZnO} = (\text{R}(\text{OC}_2\text{H}_5)_2\text{S} + \text{ZnCl}_2 + \text{H}_2\text{O})$. However, this sapon. theory is here untenable as it does not explain the formation of polymers, III, IV, R_2S , II, etc. There are here many other reactions closely connected with intermediary formation of III which by addn. of H_2O or alc. at the double bond is converted to other compds. $(\text{CH}_3\text{CH}_2)_2\text{SCH}_2$ easily takes up H_2O and $\text{R}(\text{OH})$ with formation of $(\text{HO}(\text{C}_2\text{H}_5)_2\text{SO}_2)$ and $(\text{R}(\text{OC}_2\text{H}_5)_2\text{SO}_2)$. The portion of the product of the interaction of I and Zn which is non-volatile with steam contains II, $\text{C}_2\text{H}_5\text{OR}$ (V), $(\text{HO}(\text{C}_2\text{H}_5)_2\text{S})$ (VI) and $(\text{CH}_3\text{SCH}_2\text{CH}_2\text{OR})$ (VII) and some others. The first 2 are formed thus: $\text{III} + 2\text{H}_2\text{O} = \text{VI}$, and $\text{III} + \text{H}_2\text{O} + \text{R}(\text{OH}) = \text{HO}(\text{C}_2\text{H}_5)_2\text{S}$, OR . The formation of VII is difficult to explain, and may proceed thus: $\text{I} + (\text{R}(\text{OC}_2\text{H}_5)_2\text{S}) \rightleftharpoons (\text{C}_2\text{H}_5\text{SCH}_2\text{CH}_2\text{OR}) + \text{C}_2\text{H}_5\text{OR}$. Among some other products is formed a disulfide: $2\text{HO}(\text{C}_2\text{H}_5)_2\text{S} = \text{H}_2\text{S} + (\text{C}_2\text{H}_5)_2\text{S}$. *Exptl. part.*—The procedure adopted for all $\text{R}(\text{OC}_2\text{H}_5)_2\text{SC}_2\text{H}_4\text{SC}_2\text{H}_4\text{OR}$ (VIII). To 200 g. of I in 600 cc. of 90% alc. was added 150 g. of Zn dust, this was brought to boil on a water bath, the flame removed and the acid gases were absorbed in 5% NaOH, the unsatd. gaseous compds. led into lit. water, and the unabsorbed gases were collected in a gasometer; after 1 hr. the reaction mass was refluxed 40 min. There were collected, besides C_2H_6 , 16 l. (at 20°) of unabsorbed gases, while the flask with NaOH contained Na_2S , NaCl and a little (CH_3SH) , and that with lit. water had 20 g. of $\text{C}_2\text{H}_5\text{Br}$. The main reaction mass was filtered, most of the alc. expelled from the filtrate, and then steam-distd., producing 120 g. of a volatile product, 0.5 of which is extractable with CHCl_3 while the other half is an oil lighter than H_2O and is sep'd. in the funnel. Heating 6 hrs. of I and Zn with dil. alc. (75%) reduces the volatile portion of the reaction product to 50 g. On redistn. of 500 g. of the volatile portion are obtained 5.8% b. up to 200°, 8.41% b. 200–19°, 10.60% b. 210–23°, 50.00% b. 227–7°, and 13.88% b. 227–32°. The fraction b. up to 200° on redistn. gave fractions b. 93–5° and 150–155°; the 93–5° fraction is Et_2S , d. 0.8400, gives 2(RtS) 311g. Cl_2 , m. 110°, and Rt_2S 2(HgCl), m. 127°. The fraction b. 150–155°, d. 1.1040 is vinyl

β -hydroxyethyl sulfide (IX), which gives $C_4H_{10}OS.HCl$, m. 170° , with Br. C_4H_9OSBr . m. 89° (decompos.) and with HI produces $CH_3CH_2SCH_2CH_3$, an oil with m.p. $+22.6-5^\circ$, d. 0.9672. The fraction b. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction c. $223-7^\circ$ on redistn. contains mainly a portion b. $225-65^\circ$, d. 0.9672. The fraction d. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction e. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction f. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction g. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction h. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction i. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction j. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction k. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction l. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction m. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction n. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction o. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction p. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction q. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction r. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction s. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction t. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction u. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction v. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction w. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction x. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction y. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672. The fraction z. $223-7^\circ$ on redistn. consists mainly of a portion b. $225-65^\circ$, d. 0.9672.

[illegible]

ing 36 g. of a viscous oil, d. 1.1840, which on distn. in vacuum is completely decompd. with formation of II, phenol and other compds. The oil on investigation seems to be a mixt. of $\text{PhOCH}_2\text{CH}_2\text{SCH}_2\text{CH}_2\text{OPh}$ and $\text{PhOCH}_2\text{CH}_2\text{SCH}_2\text{CH}_2\text{SCH}_2\text{CH}_2\text{OPh}$, which cannot be sep'd. by distn. On heating it with HI is obtained a mixt. of the corresponding iodides of both sulfides, while oxidation with H_2O_2 produces a liquid and crystals, both insol. in common solvents. The action of Zn, Fe and Al and their oxides could be investigated only in solvents inert to I, such as C_6H_6 , PhMe , etc. On refluxing a mixt. of 50 g. Zn dust, 50 g. of I and 100 cc. of xylene for 4 hrs., there are obtained 6 l. of gaseous matter (70-85% C_2H_4 , 25% H_2S and 3H_2), and a mass which on steam distn. produces 3 g. of unsatd. sulfides, 2 g. of II, 3.4 g. of high-mol. sulfides and polymers of II. Fine Fe and Al react similarly, the reaction proceeding with metals in neutral solvents at a higher temp. (110-200°) than in alc. or aq. alc. The secondary alcs react with I in presence of Zn dust similarly but less energetically. iso-PrOH gives $(\text{CH}_3\text{CH}_2\text{OPr})_2\text{S}$, b. 231.5-4.5°, d. 0.839; $(\text{CH}_3\text{CH}_2\text{OPr})_2\text{S} \cdot 2\text{HgCl}_2$, m. 90°. The tertiary alcs either do not react with I or react very slowly. *Conclusions.*—The action of Zn dust on I in the presence of primary and secondary fatty alcs. and of primary fatty-aromatic alcs. produces as principal products of reaction thio ethers of the types $(\text{ROCH}_2\text{CH}_2)_2\text{S}$ and $(\text{CH}_3\text{SCH}_2\text{CH}_2\text{OR})_2$. The interaction of I and phenols in the presence of Zn dust results also in thio ethers, the compn. and properties of which were not definitely detd. As by-products are formed thiodiglycol and monothethers of the formula $\text{HOCH}_2\text{CH}_2\text{SCH}_2\text{CH}_2\text{OR}$, thiodiglycol and II, IV, $(\text{CH}_3\text{SH})_2$, Et_2S , $\text{C}_6\text{H}_5\text{SH}$, H_2S , H_2 , HCl , etc. The formation of thiodiglycol and its ethers proceeds mainly by splitting off HCl from I with formation of III as an intermediary product. The latter, formed *in statu nascendi*, adds on H_2O and alc. and is converted either to ethers or thiodiglycol. A partial formation of ethers by action of Zn

10

Process and properties of sulfones and sulfoxes of the aliphatic series. Divinyl sulfone, vinyl β -chloroethyl sulfone and their derivatives. A. R. KANTOV, *J. Russ. Phys. Chem. Soc.* 62, 1: 29 (1930). The investigation of the action of Zn dust on $(\text{ClCH}_2\text{CH}_2)_2\text{S}$ (I) in alc. (C. A. 24, 3750) was extended to analogous reaction with halogen deriva. of sulfones and sulfoxides. It was found that the cyclic formation proceeds more easily with the sulfones, next in order with sulfoxides, and finally with sulfides (Cashmore, C. A. 17, 3157; Fromm and Ungar, C. A. 18, 1112; Clarke, C. A. 7, 337, 583; Arndt and Bielech, C. A. 18, 1115). The greater no. of reactions proceed without cyclization and more energetically with halogen deriva. of sulfides and sulfones. *Exptl. part.*—Divinyl sulfone, $(\text{CH}_2=\text{CH})_2\text{SO}_2$ (II): Refluxing a mixt. of 200 g. of $(\text{ClCH}_2\text{CH}_2)_2\text{SO}_2$ (III), 200 g. of Zn dust and 600 cc. 96% EtOH for 30 hrs., dilg. with 3 vols. of H_2O , extg. the sepd. oil with CHCl_3 , washing, drying and distg., yield 85 g. II, d_4^{20} 1.1790, b_p 118–21. Bis(α,β -dibromoethyl) sulfone (IV): Adding a slight excess of 4 mols. of Br to 5 g. of II, allowing to stand 24 hrs. and recrystg. 3 times from EtOH yield 7 g. IV, m. 134–5°. Bis(β -iodoethyl) sulfone (V): Refluxing 1 hr. 1.8 g. of II with 8 g. 50% HI yields 4 g. V, m. 205°; also obtained by 10 hrs. refluxing of 3 g. of III with 25 g. of NaI in 100 cc. 96% EtOH, yield 80%, m. 204°. Bis(β -bromoethyl) sulfone (VI): Refluxing 1 hr. 5 g. of II with 20 cc. of 50% HBr yields 6.5 g. VI, twice recrystd. from EtOH, m. 108–9°. Also obtained by refluxing 4 hrs. 5 g. of III with 50% HBr, yield 5.1 g., m. 112°. VI gives with PhSNa bis(β -phenylmercaptoethyl) sulfone (VIIa), m. 104°. Vinyl β -chloroethyl sulfone (VII): Refluxing 12 hrs. 200 g. of III, 200 g. of Zn dust and 600 cc. of 96% EtOH, filtering, dilg. with 2 vols. of H_2O , extg. with CHCl_3 , drying with CaCl_2 and distg., yield 120 g. of a crude VII, 40% pure VII, b_p 155–6°, d_4^{20} 1.2690. Vinyl β -iodoethyl sulfone (VIII): 12 hrs. refluxing of 4 g. of VII, 12 g. of KI and 80 cc. of 96% EtOH, expelling the EtOH, dilg. with H_2O , taking up in EtOH, drying and distg. off the EtOH yield 4.8 g. VIII, recrystd. from EtOH, m. 41°. α,β -Dibromoethyl β -chloroethyl sulfone (IX): Adding 45 g. of Br to 30 g. of VII gives IX, after 24 hrs. standing the whole is solidified and recrystd. from EtOH, m. 62°. Refluxing 1.5 hrs. 8 g. of VII with 15 g. of 50% HI gives 8.1 g. V, m. 205°. Refluxing 50 min. 4 g. of VII with 20 cc. of HBr yields 4.94 g. VI, recrystd. from EtOH, m. 112°. β -Bromoethyl β -iodoethyl sulfone (XII): Refluxing 40 min. 1.72 g. of VIII with 12 g. of HBr satd at 0° and filtering cold yield 2.5 g. XII, recrystd. from EtOH m. 116°, with PhSNa gives VIIa, m. 104–5°. β -Chloroethyl β -iodoethyl sulfone (XIII)

was prepd. in a manner similar to XII. β -Chloroethyl β -bromoethyl sulfone (XIV), m. 64° , together with VI, is formed on heating III with IIIr. α,β -Dibromoethyl β -iodoethyl sulfone (XV) and α,β,β' -tribromodiethyl sulfone (XVI) are obtained by inter-
action of VIII and Br; the mixt. of both halides is difficult to sep. α,β -Diodoethyl vinyl sulfone (XVII): To an emulsion of II and NaOH (an excess of 10%) is slowly added 2 N I in KI to permanent coloration, after standing for 2 hrs. the reaction mass is acidified, left overnight, filtered, pressed between filter paper and freed from the last traces of I in a vacuum desiccator, black crystals are obtained, having no sharp m. p., easily sol. in many org. solvents. Direct chlorination of II and VII produces a heavy mass, from which were isolated, resp., bis(α,β -dichloroethyl) sulfone (XVIII) and α,β,β' -trichloroethyl sulfone (XIX), both compds. are viscous liquids easily decomposed. II was also prepd. by refluxing a mixt. of 15 g. of III, 75 g. of ZnO and 300 cc. of 90% EtOH, dig. with 2 vols. of H_2O , extg. 3 times with $CHCl_3$, expelling the $CHCl_3$, yield 70 g., b_m 110-25°, which on redistn. gives 30 g., b_m 117-20°. VII was also obtained by refluxing a mixt. of 100 g. of III, 50 g. of ZnO and 150 cc. of 92% EtOH, yield 20 g., b_m 160°. H_2SO_4 and especially its acid salts easily combine with II and VII with 100% yield: $(CH_3CH_2)_2SO_4 + 2NaHSO_4 = (NaO)_2SCH_2CH_2SO_3Na$ (XX). $CH_3CH_2SO_3Na + CH_3CH_2Cl + 2NaHSO_4 = NaCl + NaO_2SCH_2CH_2SO_3CH_2CH_2SO_3Na$ (XXI). XXI is prepd. by refluxing 1.8 g. of II and 18 cc. of 40% NaHSO₄ for 1 hr., yield 3 g., purified by pptn. with EtOH. XXI is obtained as above in 3.2 g. yield from 2 g. of VII and 20 g. of 40% NaHSO₄. II and VII combine with H_2O , giving 100% of diglycol sulfone which in the presence of alkalis is converted to the thioxane: $(CH_3CH_2)_2SO_4 + 2H_2O = (HOCH_2CH_2)_2SO_4$ and $SO_2 \begin{matrix} CH_3CH_2 \\ \diagup \quad \diagdown \\ O \end{matrix} + H_2O$. $CH_3CH_2SO_3CH_2CH_2Cl + KOH$

$+ H_2O = KCl + (HOCH_2CH_2)_2SO_4$ and thioxane. A mixt. of 4.2 g. of II and 40 cc. of 2 N NaOH was refluxed 1 hr., cooled, filtered, the filtrate evapd. and extd. with $CHCl_3$, yield 3.7 g., needles, m. 129-30°. A mixt. of 4.9 g. of VII and 50 cc. of 2 N NaOH, treated as above, yields 4.5 g., needles, m. 120-30°. Bis(β -ethoxyethyl) sulfone (XXII) was prepd. by refluxing 2 hrs. a mixt. of 18 g. of VII in EtOH and 5.5 g. of NaOH in EtOH, filtered, the filtrate extd. with $CHCl_3$, distd., yield 17 g., redistd., b_m 180-3°, d₂₀ 1.1282. The MeO, EtO and iso-AmO deriva. were described by Cashmore (C. A. 17, 3157) without giving adequate and sufficient data. The following method gives good yields: To 100 g. of III in 300 cc. of 92% EtOH is slowly added a soln. of 42 g. NaOH in 250 cc. of EtOH, refluxed 1 hr., EtOH distd. off and XXII is extd. with Et₂O, yield 80%, b_m 175-85°. The MeO deriv. is obtained in a similar manner. For prepn. of deriva.

CO
10
Derivatives of phenacyl sulfides and their properties. A. E. Krasov, A. N. Panchenko and A. Komolovskii. *J. Gen. Chem. (U.S.S.R.)* 1, 393-400 (1931).
Bisdimethylphenacyl sulfide (I), obtained in poor yield by slowly adding with cooling and shaking 16 g. of cryst. Na₂S in 50 cc. of alc. to 25 g. of Me₂C₆H₄COCH₂Cl in 75 cc. of alc. to a faint alk. reaction, then adding 12 cc. of H₂O, and drying the wpt. oil over H₂SO₄, sol. in EtOH, Et₂O and AcOH, insol. in H₂O, m. 55°. *Dimethylphenacyl sulfide dioxime*, obtained in 0.8 g. yield (67.27%) by treating 1 g. of p-(Me₂C₆H₄COCH₂)₂S (II) in 15 cc. of alc. with 0.8 g. NH₂OH HCl and 2.5 g. of cryst. Na₂CO₃, then refluxing 2 hrs. in the water bath and pouring into cold water, m. 140-50°. I produced no oxime. *Monophenylhydrazones of III*, obtained in 4 g. yield (72.7%) by mixing 5.5 g. of III in 15 cc. of alc. with 8.3 g. of PhNHNH₂ in 10 cc. of alc., then refluxing 15 min. in the water bath, and dilg. with H₂O, m. 137°. *Monophenylhydrazones of I*, obtained from 1.1 g. of I and 1.5 g. of PhNHNH₂ as described above, m. 106-7°. (PhCOCH₂)₂S₂ obtained in 10 g. yield (23.8%) by adding with cooling 27 g. of Br in 10 cc. of AcOH to 27 g. of (PhCOCH₂)₂S (IV) in 25 cc. of 100% AcOH, then pouring into 100 cc. of water, washing the oil with water until it becomes cryst., m. 101°, sol. in org. solvents, insol. in H₂O. No corresponding dibromides of I and III could be obtained in pure state. *Phenacyl mercaptan (V)* and its homologs were not produced before. the common method for prepn. of mercaptans by the action of alc. NaHS on the halogen deriva. (PhCOCH₂Cl) gives IV. V was obtained in 30 g. yield (73.2%) by refluxing 20 hrs. on the water bath a mixt. of 42 g. of PhCOCH₂Cl in 170 cc. alc. and 134 g. of cryst. Na₂SO₃ (100% excess) in 45 cc. of H₂O, treating the cold reaction mass with HCl, extg. with Et₂O, drying with anhyd. Na₂SO₄, distg. off the Et₂O, repeatedly dissolving in hot alc., filtering, and collecting the wpt. oil on cooling; it is a light yellow oil with a faint odor, sol. in org. solvents, insol. in H₂O, gives with Pb salts (PhCOCH₂S)₂Pb. CHAS. BLANC

Arsino sulfides of the fatty and aromatic series. A. E. KRETOW AND A. YA. BERLIN.
J. Gen. Chem. (U. S. S. R.), 1, 411-8 (1931).—Bayer (Ann. 107, 279 (1858)) and Schulte
 (Ber. 13, 1955 (1882)), and later Nametkin and Nekrasov (C. A. 23, 3874) described the
 prepn. of org. arsino sulfides by the action of H_2S on arsino chlorides or the oxides. A
 series of fatty and aromatic arsino sulfides obtained by K. and B. are either new compds.
 or were prep'd. by novel methods. The results of the investigation fully support the
 findings of Blicke and Smith (C. A. 24, 4092) that the org. arsino sulfides of the formula
 $(RAs)_2S$ are monomers with normal mol. wt., while those of the formula RA_3S are
 polymers with abnormal mol. wt. The action of KSH and Na_2S on arsino chlorides in
 β -chlorovinylarsino
 produces identical compounds, with those obtained with H_2S into 2 g. of β -
 β -chlorovinylarsino
 $CH_3CH=CHAsCl_2$ yields $CH_3CH=CHAsS_2$, then bringing the reaction mass to a boil and quickly cooling.
 $CH_3CH=CHAsCl_2$ in abs. alc., then bringing the reaction mass to a boil and quickly cooling.
 $CH_3CH=CHAsCl_2$ in abs. alc., then bringing the reaction mass to a boil and quickly cooling.
 to light yellow plates with repulsive odor, sol. in CS_2 and $CHCl_3$, poorly sol. in alc. β -
 hydroxyvinylarsino sulfide, $HOCH=CHAsS_2$, obtained in 0.1 g. yield by passing H_2S
 into 2 g. of β - $HOCH=CHAsCl_2$ in alc. in the presence of $NaCO_3$, filtering, adding H_2O to
 formation of a turbidity, allowing to stand 24 hrs., sepd. the crystals from the viscous
 ppt. and washing with a little Me_2CO , m. 60-70°. β -Acetoxyethylarsino sulfide, $AcOCH_2CH_2AsS_2$, sol. in
 ppt. and washing with a little Me_2CO , m. 60-70°. β -Acetoxyethylarsino sulfide, $AcOCH_2CH_2AsS_2$, sol. in
 CH_3AsCl_2 prep'd. from 2 g. of $AcOCH_2CH_2AsCl_2$ in alc. β -Acetoxyethylarsino sulfide was prep'd. by then by a different
 method (cf. Tolman, C. A. 9, 15, and Washburn, C. A. 12, 850). By the interaction of
 H_2S and 65 g. of $EtAsCl_2$ in alc. was formed an oil, which was sepd. and dissolved in
 $CHCl_3$, then dried over CaH_2 and evap'd. in vacuo, giving 45 g. yellow oil with offensive
 odor, sol. in CS_2 and $CHCl_3$, d. 1.218, mol. wt. found 800, calcd. 810. Phenylarsino

sulfide, obtained in 78 g. yield from 66 g. of PhAsCl_2 in alc. and H_2S , m. 153° , mol. wt. found 559.1, calcd. 552. *p-Cyanophenylarsino sulfide*, obtained in 1.55-g. yield from 2 g. of $\text{p-NCC}_6\text{H}_4\text{AsCl}_2$ in alc. and H_2S , amorphous yellow ppt. sol. only in aq. NH_3 , m. 152° , mol. wt. found 435.2, calcd. 209 and 418. *p-Phenylenedisarsino sulfide*, m. 152° , mol. wt. found 435.2, calcd. 209 and 418. *p-Phenylenedisarsino sulfide*, obtained in 1.57 g. as above from 2 g. of $\text{p-C}_6\text{H}_4(\text{AsCl}_2)_2$ and H_2S , insol. in common solvents, m. $275-6^\circ$. *p-Naphthylarsino sulfide*, obtained in 1.7 g. yield from 2 g. of $\text{p-C}_{10}\text{H}_7(\text{AsCl}_2)_2$, m. 154° , mol. wt. found 455.7, calcd. 234 and 468. *p-Chlorophenylarsino sulfide*, obtained in 1.55-g. yield, m. 135° , mol. wt. found 640.3, calcd. 218 and 454. *p-Nitrophenylarsino sulfide*, obtained in 1.66-g. yield from 2 g. of $\text{p-ON-C}_6\text{H}_4(\text{AsCl}_2)_2$, yellow crystals from PhNH_2 , m. 210° , mol. wt. found 460.9, calcd. 229 and 458. *m-Nitrophenylarsino sulfide*, obtained in 0.64 g. yield from 1 g. of $\text{m-ON-C}_6\text{H}_4(\text{AsCl}_2)_2$, yellow powder, sol. in pyridine and CHCl_3 , m. $98-101^\circ$, mol. wt. found 470.3, calcd. 229 and 458. *p-Ethoxyphenylarsino sulfide*, obtained in 1.5 g. yield from 2 g. of $\text{EtOC}_6\text{H}_4(\text{AsCl}_2)_2$, m. $127-9^\circ$, sol. in C_6H_6 , Me_2CO , CHCl_3 , insol. in MeOH and EtOH . *Tetraphenyldiarsino sulfide*, $\text{Ph}_2\text{AsSAsPh}_2$ (Michaelis, Ann. 321, 145 (1902)), obtained in 1.5-g. yield from 2 g. of PhAsCl_2 , m. 64° , sol. in CHCl_3 , CS_2 and C_6H_6 . For the prepn. of arsino sulfides by the action of KHS , the corresponding arsino chlorides in alc. are treated with KHS in alc., the ppt. is dissolved in an excess of the reagent and pptd. again with dil. HCl , filtered, washed with alc. and H_2O and dried *in vacuo*. Identical products are obtained from arsino chlorides in alc. by the action of Na_2S in dil. alc. or in H_2O . CHAS. BLANC

Determination of halogens in organic compounds using sodium sulphide. - A. E. KNETOV, A. N. PANTUCHENKO, and K. K. RAVITSCH (J. Gen. Chem. Russ., 1931, 4, 419-422).—0.2 g. of substance is dissolved in 5 c.c. of EtOH, 7–8 c.c. of a solution containing 0.5–0.6 g. of Na₂S·9H₂O in EtOH are added, and the mixture is heated for 30 min. under reflux at 100°, cooled, diluted with H₂O to 100 c.c., feebly acidified, and boiled until all the EtOH is expelled. Residual H₂S is oxidized by adding 3 c.c. of 30% H₂O₂ to the alkaline solution, which is then boiled a few min. to destroy excess of H₂O₂, and ionic halide is determined by Volhard's method. When coloured solutions are obtained after the action of H₂O₂, they should be further oxidized by KMnO₄ to the disappearance of the coloration. The above method is applicable to halogens present in aliphatic but not aromatic groups; when they are present in both the nucleus and the side-chain of aralkyl compounds only those of the side-chain are determined.
R. TRUSZKOWSKI.

Oxidizing properties of halonitromethanes. A. R. KERRIOV AND N. M. MALHIKOV
J. Gen. Chem. (U. S. S. R.) 2, 202-7(1932).--Nitro compds. act as strong oxidizing
agents, undergoing some intramol. oxidation with disintegration of the mol (Nef, *Ann*
280, 273(1904)), while halonitro deriva. react with even greater oxidizing energy, under-
going highly complex reactions and producing a considerable no. of decompn. products.
The oxidizing properties of chloropicrin (I) and bromopicrin (II) were further established
(cf. Ray, Guha and Das, *C. A.* 14, 1238; 16, 1933; Nekrasov and M., *C. A.* 24, 87)
by the pptn. of I in the interaction of I and KI (Shchev, *C. A.* 23, 4210) by their
reactions with KI and alkali sulfides and hydrosulfides and mercaptans. I and NaHS
Pure cryst. NaHS $2H_2O$ (27 g.) was dissolved in 250-300 cc. of KOH by refluxing, the
slr being expelled from the system by the alc. vapors; then 15.3 g. of I was introduced
drop by drop in about 1.5 hrs. while the soln. was kept boiling. In some expts. the
escaping gases were absorbed in satd. alc. KOH, while for the detn. of such gases as N_2O
and CO_2 in the miats. sep. expts. were required. Besides CO_2 , there was obtained 410
40 cc. gases contg. NO 14.0-17.2, CO_2 2.4-4.0, O_2 0.1, and N 79.2-80%. The reaction
flask contained 3.0-4.1 g. of elementary S, 21.5-5.6 g. of NaCl and 0.3-0.9 g. of $NaNO_3$.
 NH_3 and NH_4OH were detected but not detd. CS_2 absorbed in alc. KOH and converted
quantitatively to $CS(OEt)SK$ was pptd. as the Cu salt and analyzed by titrating with I
and detg. the S. I and K_2S . Under similar conditions, 22.4 g. of pure cryst. $K_2S \cdot 2H_2O$ in
250 cc. of alc. produced with I, besides CO_2 , 510-40 cc. of gases contg. NO 22.54-25,
 CO_2 30.33-7, O_2 0.7, CO 4-8 and N 33.1-35.7%. No NH_3 and NH_4OH were found.
A salt cake (10.8-20 g.) was formed contg. mostly KCl, 0.4 g. of KNO_3 and 0.5 g. S. In
the alc. KOH was found considerable $EtOCSOK$ formed by the interaction of CO_2 and
alc. KOH. No CS_2 was found in the alc. KOH. II and Na_2S . To 90 g. of $Na_2S \cdot 9H_2O$
in 250 cc. of alc. was slowly added 75 g. of II in 75 cc. of alc., and then the mixt. was
heated 15-30 min. to complete the reaction. The gas contained CO_2 30.5, NO 12.4.

***β,β'*-Dichloro- and *β,β'*-dibromodivinyl sulfoxides, and their properties.** A. E. KARTOV AND B. M. KLIGER. *J. Gen. Chem. (U. S. S. R.)* 2, 322-6 (1932).—The study of halogen derivs. of $S(CH_2CH_2Cl)_2$ and its products of oxidation led to synthesis and investigation of a large no. of 8 compds. (Lawson and Dawson, *C. A.* 22, 381; Mumford and Phillips, *C. A.* 22, 1325; Phillips, Davies and Mumford, *C. A.* 23, 2932; Mann and Poppe, *C. A.* 16, 2110). Some of the most interesting and least available compds. of this type are halo sulfoxides with double bonds and their products of oxidation. $OS(CH_2CH_2Cl)_2$ (I) and $CH_2=CHSOCH_2CH_2Cl$ (II) were obtained by treating $OS(CH_2CH_2Cl)_2$ (III) with $NaHCO_3$; $III + NaHCO_3 \rightarrow II + H_2O + CO_2 + NaCl$, and $III + 2NaHCO_3 \rightarrow I + 2H_2O + 2CO_2 + 2NaCl$. By chlorination and bromination of I are obtained, resp., *α,α',β,β'*-tetrachlorodivinyl sulfoxide (IV) and *α,α',β,β'*-tetrabromodivinyl sulfoxide (V), which by splitting off 1 hydrogen halide with $NaHCO_3$ give *β,β'*-dichlorodivinyl sulfoxide (VI) and *β,β'*-dibromodivinyl sulfoxide (VII). The failure of VI and VII to produce with H the corresponding sulfides, and to give some reactions analogous to those of sulfoxides which have no halogen at the C united by the double bond, may be explained by the presence of 2 vinyl groups and 2 atoms of halogen in the *β*-positions. *Exptl. part.* IV, m. 122.5°, was prepd. in 85% yield by conducting 15.4 g. of dry Cl (110% of theory obtained from the calcd. amt. of pure $KMnO_4$ and an excess of HCl) into 10 g. of I in 100 cc. of ice-cold abs. alc.; at the end of 3 hrs the temp. rose to 20°; the mixt. was allowed to stand overnight, and the white crystals were filtered off. The oxidation of 5 g. IV with $KMnO_4$ and H_2SO_4 (Müller and Metzger, *C. A.* 21, 52) gave 1 g. of $ClCH_2CHClSO_2K$. VI, b.p. 86-7°, m. 15°, d_4^{20} 1.417, was obtained in 85% yield by blowing steam into a mixt. of 64 g. of IV and 43 g. of $NaHCO_3$ in water, extg. the steam distillate with Et_2O , drying with anhyd. Na_2SO_4 , and fractionating. V, m. 120.8°, was prepd. in 35-50% yield like IV by conducting gaseous Br into I. VII, m. 40.5-2°, was obtained in 75% yield as VI. CHAS. BLANC

cc

18

Different methods of production of cyanide compounds. A. B. Kretov, J. Chem.
Ind. (Moscow) 1933, No. 4, 28-34. Review.

H. M. Leicester

ASH-SLA METALLURGICAL LITERATURE CLASSIFICATION

10

✓ This ketone of aliphatic series. I. The action of phosphorus pentasulfide on aliphatic ketones. A. E. Kretov and Ya. F. Kondimarov. *J. Gen. Chem.* (U. S. S. R.) 8, 255-51 (1935).—Mixts. of powd. P_2S_5 with Me_2CO , $MeCOEt$, Et_2CO , Pr_2CO and $(Me_2C)_2CO$ in PhMe were refluxed on a water bath for 8 hrs., the reaction mixts. were steam-distd., the solns. of the thio ketones in PhMe were dried with $CaCl_2$, the PhMe distd. off *in vacuo* and the residues fractionated. Most of the thio ketones were obtained in small yields and impure state, boiling *in vacuo* within 10° . They are yellow-red liquids somewhat heavier than H_2O , easily sol. in org. solvents. Me_2CS , $MeCSEt$ and Et_2CS are dimers, Pr_2CS a mixt. of a monomer with some dimer and $(Me_2C)_2CS$ a monomer. (Me_2CS)₂, b. $182-5^\circ$. ($MeCSEt$)₂, b. $120-30^\circ$, mol. wt. 167 (found), 170 (calcd.), gives ($MeCSEt$)₂, $HgCl_2$. (Et_2CS)₂, b. $135-50^\circ$, mol. wt. 210 (found), 204 (calcd.), gives the $HgCl_2$ compd. Pr_2CS , b. $135-60^\circ$ (slight decompn.), mol. wt. 169 and 173 (found), 130 (calcd.). (Me_2C)₂CS, b. $105-10^\circ$, gives the $HgCl_2$ compd. C. B.

ASM-SLA METALLURGICAL LITERATURE CLASSIFICATION

137 AND 138 (EXPT)

PROCESSES AND PROPERTIES INDEX

139 AND 140 (EXPT)

141 AND 142 (EXPT)

143 AND 144 (EXPT)

145 AND 146 (EXPT)

147 AND 148 (EXPT)

149 AND 150 (EXPT)

151 AND 152 (EXPT)

153 AND 154 (EXPT)

155 AND 156 (EXPT)

157 AND 158 (EXPT)

159 AND 160 (EXPT)

161 AND 162 (EXPT)

163 AND 164 (EXPT)

165 AND 166 (EXPT)

167 AND 168 (EXPT)

169 AND 170 (EXPT)

171 AND 172 (EXPT)

173 AND 174 (EXPT)

175 AND 176 (EXPT)

177 AND 178 (EXPT)

179 AND 180 (EXPT)

181 AND 182 (EXPT)

183 AND 184 (EXPT)

185 AND 186 (EXPT)

187 AND 188 (EXPT)

189 AND 190 (EXPT)

191 AND 192 (EXPT)

193 AND 194 (EXPT)

195 AND 196 (EXPT)

197 AND 198 (EXPT)

199 AND 200 (EXPT)

201 AND 202 (EXPT)

203 AND 204 (EXPT)

205 AND 206 (EXPT)

207 AND 208 (EXPT)

209 AND 210 (EXPT)

211 AND 212 (EXPT)

213 AND 214 (EXPT)

215 AND 216 (EXPT)

217 AND 218 (EXPT)

219 AND 220 (EXPT)

221 AND 222 (EXPT)

223 AND 224 (EXPT)

225 AND 226 (EXPT)

227 AND 228 (EXPT)

229 AND 230 (EXPT)

231 AND 232 (EXPT)

233 AND 234 (EXPT)

235 AND 236 (EXPT)

237 AND 238 (EXPT)

239 AND 240 (EXPT)

241 AND 242 (EXPT)

243 AND 244 (EXPT)

245 AND 246 (EXPT)

247 AND 248 (EXPT)

249 AND 250 (EXPT)

251 AND 252 (EXPT)

253 AND 254 (EXPT)

255 AND 256 (EXPT)

257 AND 258 (EXPT)

259 AND 260 (EXPT)

261 AND 262 (EXPT)

263 AND 264 (EXPT)

265 AND 266 (EXPT)

267 AND 268 (EXPT)

269 AND 270 (EXPT)

271 AND 272 (EXPT)

273 AND 274 (EXPT)

275 AND 276 (EXPT)

277 AND 278 (EXPT)

279 AND 280 (EXPT)

281 AND 282 (EXPT)

283 AND 284 (EXPT)

285 AND 286 (EXPT)

287 AND 288 (EXPT)

289 AND 290 (EXPT)

291 AND 292 (EXPT)

293 AND 294 (EXPT)

295 AND 296 (EXPT)

297 AND 298 (EXPT)

299 AND 300 (EXPT)

301 AND 302 (EXPT)

303 AND 304 (EXPT)

305 AND 306 (EXPT)

307 AND 308 (EXPT)

309 AND 310 (EXPT)

311 AND 312 (EXPT)

313 AND 314 (EXPT)

315 AND 316 (EXPT)

317 AND 318 (EXPT)

319 AND 320 (EXPT)

321 AND 322 (EXPT)

323 AND 324 (EXPT)

325 AND 326 (EXPT)

327 AND 328 (EXPT)

329 AND 330 (EXPT)

331 AND 332 (EXPT)

333 AND 334 (EXPT)

335 AND 336 (EXPT)

337 AND 338 (EXPT)

339 AND 340 (EXPT)

341 AND 342 (EXPT)

343 AND 344 (EXPT)

345 AND 346 (EXPT)

347 AND 348 (EXPT)

349 AND 350 (EXPT)

351 AND 352 (EXPT)

353 AND 354 (EXPT)

355 AND 356 (EXPT)

357 AND 358 (EXPT)

359 AND 360 (EXPT)

361 AND 362 (EXPT)

363 AND 364 (EXPT)

365 AND 366 (EXPT)

367 AND 368 (EXPT)

369 AND 370 (EXPT)

371 AND 372 (EXPT)

373 AND 374 (EXPT)

375 AND 376 (EXPT)

377 AND 378 (EXPT)

379 AND 380 (EXPT)

381 AND 382 (EXPT)

383 AND 384 (EXPT)

385 AND 386 (EXPT)

387 AND 388 (EXPT)

389 AND 390 (EXPT)

391 AND 392 (EXPT)

393 AND 394 (EXPT)

395 AND 396 (EXPT)

397 AND 398 (EXPT)

399 AND 400 (EXPT)

401 AND 402 (EXPT)

403 AND 404 (EXPT)

405 AND 406 (EXPT)

407 AND 408 (EXPT)

409 AND 410 (EXPT)

411 AND 412 (EXPT)

413 AND 414 (EXPT)

415 AND 416 (EXPT)

417 AND 418 (EXPT)

419 AND 420 (EXPT)

421 AND 422 (EXPT)

423 AND 424 (EXPT)

425 AND 426 (EXPT)

427 AND 428 (EXPT)

429 AND 430 (EXPT)

431 AND 432 (EXPT)

433 AND 434 (EXPT)

435 AND 436 (EXPT)

437 AND 438 (EXPT)

439 AND 440 (EXPT)

441 AND 442 (EXPT)

443 AND 444 (EXPT)

445 AND 446 (EXPT)

447 AND 448 (EXPT)

449 AND 450 (EXPT)

451 AND 452 (EXPT)

453 AND 454 (EXPT)

455 AND 456 (EXPT)

457 AND 458 (EXPT)

459 AND 460 (EXPT)

461 AND 462 (EXPT)

463 AND 464 (EXPT)

465 AND 466 (EXPT)

467 AND 468 (EXPT)

469 AND 470 (EXPT)

471 AND 472 (EXPT)

473 AND 474 (EXPT)

475 AND 476 (EXPT)

477 AND 478 (EXPT)

479 AND 480 (EXPT)

481 AND 482 (EXPT)

483 AND 484 (EXPT)

485 AND 486 (EXPT)

487 AND 488 (EXPT)

489 AND 490 (EXPT)

491 AND 492 (EXPT)

493 AND 494 (EXPT)

495 AND 496 (EXPT)

497 AND 498 (EXPT)

499 AND 500 (EXPT)

501 AND 502 (EXPT)

503 AND 504 (EXPT)

505 AND 506 (EXPT)

507 AND 508 (EXPT)

509 AND 510 (EXPT)

511 AND 512 (EXPT)

513 AND 514 (EXPT)

515 AND 516 (EXPT)

517 AND 518 (EXPT)

519 AND 520 (EXPT)

521 AND 522 (EXPT)

523 AND 524 (EXPT)

525 AND 526 (EXPT)

527 AND 528 (EXPT)

529 AND 530 (EXPT)

531 AND 532 (EXPT)

533 AND 534 (EXPT)

535 AND 536 (EXPT)

537 AND 538 (EXPT)

539 AND 540 (EXPT)

541 AND 542 (EXPT)

543 AND 544 (EXPT)

545 AND 546 (EXPT)

547 AND 548 (EXPT)

549 AND 550 (EXPT)

551 AND 552 (EXPT)

553 AND 554 (EXPT)

555 AND 556 (EXPT)

557 AND 558 (EXPT)

559 AND 560 (EXPT)

561 AND 562 (EXPT)

563 AND 564 (EXPT)

565 AND 566 (EXPT)

567 AND 568 (EXPT)

569 AND 570 (EXPT)

571 AND 572 (EXPT)

573 AND 574 (EXPT)

575 AND 576 (EXPT)

577 AND 578 (EXPT)

579 AND 580 (EXPT)

581 AND 582 (EXPT)

583 AND 584 (EXPT)

585 AND 586 (EXPT)

587 AND 588 (EXPT)

589 AND 590 (EXPT)

591 AND 592 (EXPT)

593 AND 594 (EXPT)

595 AND 596 (EXPT)

597 AND 598 (EXPT)

599 AND 600 (EXPT)

601 AND 602 (EXPT)

603 AND 604 (EXPT)

605 AND 606 (EXPT)

607 AND 608 (EXPT)

609 AND 610 (EXPT)

611 AND 612 (EXPT)

613 AND 614 (EXPT)

615 AND 616 (EXPT)

617 AND 618 (EXPT)

619 AND 620 (EXPT)

621 AND 622 (EXPT)

623 AND 624 (EXPT)

625 AND 626 (EXPT)

627 AND 628 (EXPT)

629 AND 630 (EXPT)

631 AND 632 (EXPT)

633 AND 634 (EXPT)

635 AND 636 (EXPT)

637 AND 638 (EXPT)

639 AND 640 (EXPT)

641 AND 642 (EXPT)

643 AND 644 (EXPT)

645 AND 646 (EXPT)

64

17

ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION

ANABENINE SULFONATE. A. G. Sokolov and A. E. Kretov. Rum. 61,137, May 31, 1937. A soln. of anabrine or its sulfate is treated consecutively with H_2SiF_6 and EtOH or MeOH.

15

Some tetraarylophosphonium chlorides. N. N. Mel'nikov, A. B. Kretov and B. I. Mel'tser. *J. Gen. Chem.* (U. S. S. R.) 7, 461-3 (1937).—Halogen-substituted ketones or benzyl derivs. react with Ph_4P^+ to give *triphenylacetonylphosphonium chloride* m. 234° (decomp.), its *bromide* m. 221°; *triphenylphenacylphosphonium bromide* m. 253° (decomp.); *triphenyl o-nitrobenzylphosphonium chloride* m. 231° (decomp.), its *m-isomer* m. 247° (decomp.), its *p-isomer* m. 247° (decomp.) and *triphenyl o-cyanobenzylphosphonium chloride* m. 244-5°.

! The NO_2 - and CN-substituted compds. undergo some oxidation during their prepn. H. M. Leicester

ASB-51-A METALLURGICAL LITERATURE CLASSIFICATION

SECTION	SUBSECTION	CLASSIFICATION
1	1	1
2	2	2
3	3	3
4	4	4
5	5	5
6	6	6
7	7	7
8	8	8
9	9	9
10	10	10
11	11	11
12	12	12
13	13	13
14	14	14
15	15	15
16	16	16
17	17	17
18	18	18
19	19	19
20	20	20
21	21	21
22	22	22
23	23	23
24	24	24
25	25	25
26	26	26
27	27	27
28	28	28
29	29	29
30	30	30
31	31	31
32	32	32
33	33	33
34	34	34
35	35	35
36	36	36
37	37	37
38	38	38
39	39	39
40	40	40
41	41	41
42	42	42
43	43	43
44	44	44
45	45	45
46	46	46
47	47	47
48	48	48
49	49	49
50	50	50
51	51	51
52	52	52
53	53	53
54	54	54
55	55	55
56	56	56
57	57	57
58	58	58
59	59	59
60	60	60
61	61	61
62	62	62
63	63	63
64	64	64
65	65	65
66	66	66
67	67	67
68	68	68
69	69	69
70	70	70
71	71	71
72	72	72
73	73	73
74	74	74
75	75	75
76	76	76
77	77	77
78	78	78
79	79	79
80	80	80
81	81	81
82	82	82
83	83	83
84	84	84
85	85	85
86	86	86
87	87	87
88	88	88
89	89	89
90	90	90
91	91	91
92	92	92
93	93	93
94	94	94
95	95	95
96	96	96
97	97	97
98	98	98
99	99	99
100	100	100

1ST AND 2ND CODES										3RD AND 4TH CODES										5TH AND 6TH CODES																																																																					
PROCESSES AND PROPERTIES INDEX																																																																																									
BC																				AB																																																																					
<p>Preparation of mercury organic compounds by direct action of metallic mercury. A. E. KAROV and V. A. ABRAMOV (J. Gen. Chem. Russ., 1937, 7, 1872-1878). --CHPhBr-CN in COMeEt and Hg (8 hr. at 40°) yield a mixture of phenylbromo- mercuriacetonitrile, m.p. 157°, and of its decomp. product, dicyanodibenzyl, m.p. 237°. R. T.</p>																																																																																									
AUG. 5.1.1 METALLURGICAL LITERATURE CLASSIFICATION																																																																																									
FROM SYNONYMS																																																																																									
SEARCHED 02										SEARCHED 03 04 05 06										COLLECTED 07																																																																					
<table border="1"> <tr> <td>01</td><td>02</td><td>03</td><td>04</td><td>05</td><td>06</td><td>07</td><td>08</td><td>09</td><td>10</td> <td>11</td><td>12</td><td>13</td><td>14</td><td>15</td><td>16</td><td>17</td><td>18</td><td>19</td><td>20</td> <td>21</td><td>22</td><td>23</td><td>24</td><td>25</td><td>26</td><td>27</td><td>28</td><td>29</td><td>30</td> </tr> <tr> <td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td> <td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td> <td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td> </tr> </table>																														01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30																														
01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30																																																												

ica

10

Thiocyanate sulfides and sulfones. A. E. Kretov and E. M. Toropova. *J. Gen. Chem.* (U. S. S. R.) 7, 2000-15 (1937).—When EtSNa is treated slowly with $\text{Cl}(\text{CH}_2)_3\text{Br}$ (I) and the mixt. is heated at $80-90^\circ$, the product is γ -chloropropyl ethyl sulfide (II) b.p. $68-72^\circ$. Similarly, PhSNa and I give $\text{PhS}(\text{CH}_2)_3\text{Cl}$ (III). Oxidation of $\text{Cl}(\text{CH}_2)_3\text{SH}$ (IV) by H_2O_2 gives $\text{Cl}(\text{CH}_2)_3\text{SO}_2\text{H}$, b.p. $120-2^\circ$, d^{20}_4 1.2830, n^{20}_D 1.4760. In a similar way is obtained $\text{Cl}(\text{CH}_2)_3\text{SO}_2\text{Ph}$, b.p. $108-74^\circ$, m. 82° . H_2O_2 and II give $\text{Cl}(\text{CH}_2)_3\text{SO}_2\text{Et}$, b.p. $100-3^\circ$, d^{20}_4 1.2451, n^{20}_D 1.4749, and also $\text{EtSO}_2(\text{CH}_2)_3\text{SO}_2\text{Et}$, m. $181-2^\circ$. III and H_2O_2 form $\text{Cl}(\text{CH}_2)_3\text{SO}_2\text{Ph}$, m. $23-4^\circ$. IV and KSCN in EtOH at $60-70^\circ$ for 1-1.5 hrs. give $\text{EtS}(\text{CH}_2)_3\text{SCN}$, b.p. $108-10^\circ$. In the same way were prepd. $\text{PhS}(\text{CH}_2)_3\text{SCN}$, b.p. $145-6^\circ$; $\text{EtS}(\text{CH}_2)_3\text{SCN}$, b.p. $115-20^\circ$, d^{20}_4 1.086, n^{20}_D 1.5230; and $\text{PhS}(\text{CH}_2)_3\text{SCN}$, b.p. $176-8^\circ$, d^{20}_4 1.160, n^{20}_D 1.587. Oxidation of these sulfides with H_2O_2 gives the corresponding sulfones: $\text{KSH}(\text{CH}_2)_3\text{SCN}$, m. $36-7^\circ$, $\text{PhS}(\text{CH}_2)_3\text{SCN}$, m. $71.6-8^\circ$, $\text{EtSO}_2(\text{CH}_2)_3\text{SCN}$, m. $89.5-41^\circ$, and $\text{PhSO}_2(\text{CH}_2)_3\text{SCN}$, m. 91° . The thiocyanate compds. contg. a Ph group are more toxic to insects than the purely aliphatic compds. and the sulfides are more toxic than the sulfones. However, these compds. are less toxic than ethylene dithiocyanate or aniline thiocyanate. H. M. Leicester

Reaction of α -nitrophenylsulfur chloride with potassium hydrosulfide. Gregg Dougherty and Otto Haas. *J. Am. Chem. Soc.* 59, 2460-70 (1937). $\alpha\text{-O}_2\text{NC}_6\text{H}_4\text{SCl}$ and KSH in abs. EtOH at room temp. give $(2\text{-O}_2\text{NC}_6\text{H}_4\text{S})_2$ (I) KCl , S and H_2S . It is assumed that $\text{O}_2\text{NC}_6\text{H}_4\text{SH}$ is an intermediate product, which decomps. as rapidly as formed to I, S and H_2S . An explanation of the formation of thianthrene from $\text{C}_6\text{H}_5\text{S}$ and AlCl_3 , based on this reaction, is given. C. I. West

ASM-SLA DETAILURGICAL LITERATURE CLASSIFICATION

KRETOV, A. E.

Chemical Abstr.

Vol. 48 No. 8

Apr. 25, 1954

Fuels and Carbonization Products

Chlorination of Ukrainian brown coal. A. P. Kretov,
M. I. Savin, M. I. Shenbor, and I. I. Leschenko, ~~Ukrainian~~
~~Ukr. Dnepropetrovsk~~, *Ukrain. Khim. Zvez.* 1954, 1, 10
(1953) (in Russian).—Ukrainian brown coal is easily chlori-
nated, yielding products that are sol. in org. solvents and
are reactive. The products can be used as a source of fuel
and resins and film-forming materials. The products are
light brown to orange and contain up to 1.5% Cl.
Chlorination is possible in CCl_4 medium or in the presence
of H_2O ; in the latter case the reaction is substantially com-
plete within 10 hrs. at 0-60° temp. range. G. M. K.

9-16-54
JAP

Salts of benzidine with carboxylic acids of the aliphatic series and the application of benzidine for determination of cis-trans isomers of unsaturated dicarboxylic acids. A. E. Kretov (Dnepropetrovsk Chem. Technol. Inst.). *Sbornik Nauch. Obzrach. Khim., Akad. Nauk S.S.S.R.* 1, 510-16 (1953).—Benzidine (I) forms with trans unsatd. dicarboxylic aliphatic acids almost colorless salts, while with cis isomers of the same acids, the salts are yellow-green or yellow. This fact can be used for detn. of configuration of the acids. Thus 2.3 g. I in EtOH treated with 2 g. fumaric acid gave the salt $C_{12}H_{10}O_4N_4$, m. 310-20°; reverse addn. gave the 1:2 salt $C_{12}H_{10}O_4N_4$, with a lower m. p. Reaction of 2.4 g. I in EtOH added to 2 g. maleic acid gave yellow-green salt, $C_{12}H_{10}O_4N_4$, whose m. p. is high and not const. owing to decompn.; addn. of 2 g. maleic acid to 0.5 g. I in EtOH gave grey-yellow salt, $C_{12}H_{10}O_4N_4$, m. 312-14°. Mesaconic acid (2 g.) added to 2.3 g. I in EtOH gave colorless $C_{12}H_{10}O_4N_4$, m. 174-5°; reverse order of mixing gave the same product. Citraconic acid and I gave the yellow salt, $C_{12}H_{10}O_4N_4$, decomp. about 330°. Aconitic acid and I gave also only $C_{12}H_{10}O_4N_4$. [*p*- $C_6H_4(NH_2)_2$] added to excess maleic acid in EtOH gave a grey-blue salt, $C_{12}H_{10}O_4N_4$, decomp. 165°. With dibasic acids I formed a series of salts listed below with the molar ratio of the acid to I given in parentheses; oxalic (1:1), decomp. above 400°; parabanic (1:1), m. 149°; malonic (1:2), m. 158-0°; chloromalonic (1:1), decomp. 340°; succinic (1:2), m. 184-5°; dibromosuccinic (1:1), decomp. with loss of Br; glutaric (1:2), m. 120-1°; adipic (1:2), m. 140°; chloromaleic (1:1), m. p. not sharp; citric (1:2), decomp.; pyromucic (1:2), m. 332-4°; tricarballic (1:2), m. 161-2°; $ClCH_2CO_2H$ (2:1), m. 283-9°; tartaric (1:1), decomp.; acetylenedicarboxylic (1:1); itaconic (1:1). While aconitic acid forms a 1:3 salt with I, the unsatd. cis-isomers of dicarboxylic acids form salts with 2 moles I per 3 moles acid. Stronger acids give 1:1 salts. G. M. Kowolapoff

K. K. Fed. A.E.

USSR

Reaction of *N,N*-dichlorobenzene-sulfonamide with alcohols. I. Reaction of *N,N*-dichlorobenzene-sulfonamide with primary fatty alcohols. A. P. Kretov and K. A. Chervinskii. *Ukrain. Khim. Zh.* 1954, 10, 159-63 (1954); *Russk. Zhur., Khim.* 1954, No. 21502. The reaction of *N,N*-dichlorobenzene-sulfonamide (I) with MeOH (II), EtOH (III), BuOH (IV), iso-BuOH (V), or octyl alc. (VI) proceeds as follows: $\text{PhSO}_2\text{NCl}_2 + 2\text{RCH}_2\text{OH} \rightarrow \text{PhSO}_2\text{NH}_2 + \text{RCOCH}_2\text{R} + \text{HCl}$. As by-products are formed acetal, monochloroacetal, a chlorinated ester, and an α -chloro ether and products more highly chlorinated. The rate of reaction increases from II to VI. The reaction is accelerated by light and is autocatalytic (catalyzed by HCl); in the presence of pyridine the rate of reaction is lowered 100-fold. It is assumed that the true oxidizing agent is mol. Cl liberated during the reaction. The aldehyde formed produces a semi acetal which under the influence of Cl or HCl transforms into one of the end-products of the reaction. A soln. of 15.2 g. of I in 200 ml. of II irradiated for 8 hrs. at 20°, the mixt. neutralized with Na_2CO_3 yields upon distn. HCO_2Me 33 and methylal 18%. To a mixt. of 180 g. C_6H_5 and 15.2 g. of III is added over a period of 3 hrs. 37.7 g. of I; in the reaction mixt. was found: Et acetate 64, acetal 17.0, aldehyde 2, and combined Cl 0.4%. Upon oxidation of 24.7 ml. of IV in 160 ml. of C_6H_5 with 37.7 g. of I at 5-10° was obtained after distn. BuO_2CPr 60.5, acetal 7.8, chloroacetal 5.5, and chlorobutyl Bu ether 0.2%. Oxidation of 180 ml. V with 56.5 g. I in the presence of 50 g. pyridine (60 hrs. at approx. 20° and 2 hrs. at 50°) followed by steam distn. and fractionation of the distillate was obtained: PrCHO 18 and iso- BuO_2CPr 65%. Oxidation of 21.7 g. of VI in 175 ml. C_6H_5 with 18.8 g. of I was obtained 21.3 g. of a substance b. 80-110° in which was found:

OVER

octyl caprylate 15.1 and combined Cl 2.1%. II. Reaction of *N,N*-dichlorobenzene-sulfonamides with benzyl alcohol. K. A. Chernin'ka and A. E. Kretov. *Ukrain. Khim. Zhur.* 19, 401-4; *Referat. Zhur., Khim.* 1954, No. 21393. *N,N*-Dichlorobenzene-sulfonamide (I) reacts violently with PhCH_2OH (II). The reaction can be carried out only with small concns. of I and intense cooling. The reaction products are Benz (III) and benzyl benzoate (IV) in approx. equal yields. In C_6H_6 soln. light increases the total yield and the relative yield of IV. In CCl_4 the total yield is lower than in C_6H_6 and the yield of III somewhat higher. Light lowers the total yield at the expense of III while the yield of IV increases. It is assumed that in the decompn. of alkyl hypochlorites (V), a radical chain reaction, the 1st stage is disson. $\text{RCH}_2\text{OCl} \rightleftharpoons \text{RCH}_2\text{O}\cdot + \text{Cl}\cdot$ which occurs under the influence of light, increase of temp., or by interaction with alc. mols. In the end forms an aldehyde and HCl ($\text{RCH}_2\text{OCl} \rightarrow \text{RCHO} + \text{HCl}$ (cf. preceding abstr.)). Aldehyde forms with alc. a semi acetal which is oxidized to an ester and HCl catalyzes the reaction of I with alc. The rate of V decompn. detcs. the rate of alc.

2/3

oxidation and formation of ester in CCl_4 . V can be oxidized because of an equilibrium reaction with the solvent ($\text{V} + \text{CCl}_4 \rightleftharpoons \text{HCl} + \text{OCCl}_2 + \text{C}_6\text{H}_5$). This lowers the concentration of the aldehyde and the rate of hemiacetal formation which in turn results in a lower yield of ester. The yield of the end products can also be lowered because of chlorination reaction with atomic Cl liberated from V or I. Light hastens the decomposition of V and favors an increased yield of ester but at the same time increases also decomposition and liberation of Cl which causes an increase in chlorination products and a drop in the yield of the basic product. In C₆H₆, the decomposition of V proceeds rapidly and the chain reaction with Cl are soon arrested; this causes an increase in the yield of ester. With an excess of II (without solvent) is formed dibenzyl ether. This is formed during distillation of the reaction mixture containing an excess of II and saturated with HCl. It was shown that upon distillation of pure II after letting it stand with HCl was formed dibenzyl ether with a yield of approx. 30%. In the presence of pyridine the yield of dibenzyl ether is higher, presumably because the HCl fixed by pyridine does not volatilize in the 1st stages of distillation. Based on this it was assumed that the reaction of I with a mixture of 2 alcohols, differing widely in their rate of reaction with I will form an ester with an acid radical derived from the rapidly reacting alc. and an alc. radical derived from the slow reacting alc. Indeed, in the reaction of I with a mixture of II (explosive reaction) and MeOH (reaction requiring 7 days) was obtained Me benzoate with a yield of approx. 69.3%. This is explained as follows: first, II is oxidized to III, III forms a hemiacetal of MeOH, and this on oxidation produces Me benzoate.

M. Haseb...

IRUCOV, A. Z.

USSR.

✓ Preparation of benzylamine from benzyl chloride and
urea. A. F. Kirelov and E. D. Radchenko. *J. Appl.
Chem. U.S.S.R.* 24:206-8 (1954) Engl. translation. See
C.A. 48, 9944c.
H. L. H.

③⁶

Preparation of benzylamine from benzyl chloride and urea. A. E. Kirelov and E. N. Rudchenko (Dnepropetrovsk Chem. Technol. Inst.), *Zhur. Priklad. Khim.* 26, 1418 (1953).—In the reaction of PhCH_2Cl with urea there are first formed mono- and dibenzylurea, which, heated with alkali yield PhCH_2NH_2 . The best yield of benzyl- and dibenzylurea is 70% (based on RCl), obtained by refluxing 1 mole PhCH_2Cl with 2 moles urea and 0.83 mole H_2O 5 hrs. Generally higher temps. and lesser concns. of H_2O result in a higher dibenzylurea content in the product. The alkylated urea fused with NaOH at 250-75° yields 94-5% techn. PhCH_2NH_2 . Redistn. gives the pure amine, b. 183-5°, d₄²⁰ 0.8902, n_D²⁰ 1.5300. G. M. Kozolapoff.

NR 100, AL

U.S.S.R.

Mobile, of halogen in 1-halo derivatives of anthracene.
I. H. L. Arsen'yev and A. P. Kravtsov. J. Appl. Chem. U.S.S.R.
S.R. 46, 1219-24 (1953) (Engl. translation). H. 162.
1240-51. See C.A. 46, 18670cd. H. L. H.

PRETOV, A.E.

Mobility of halogen in 1-halo derivatives of anthraquinones.
 I. N. I. Arsenov and A. V. Krut' (Leningrad, 1953). *Zh. Fiz. Khim.* 27, 1210-1211. (1953) G. Ulmann and Minaev, *C.A.* 6, 1440. The Minaev-Ulmann reaction proceeds in the direction of formation of anthraquinone from 1-chloroanthraquinone in the presence of acetates of K or Na in polar solvents such as PhNO_2 , $1,2,4\text{-C}_6\text{H}_3\text{Cl}_3$, Ph_2O , Ph_2SO ; the K salts accelerate the reaction some 2-3 times more than do the Na salts. K_2CO_3 , Na_2CO_3 , Na silicates, etc. Instead of the org. acid salts give dianthraquinone. In cases in which anthraquinone is the principal reaction product, the by-products are: 1.8-2.0% 1-hydroxyanthraquinone and 2-2.5% dihaloanthraquinone. In nonpolar solvents the cleavage of haloanthraquinone proceeds similarly, but the reaction is much less rapid. PhNO_2 is not a specific substance in the reaction; Ph_2SO gives analogous results. $\text{C}_6\text{H}_5\text{Cl}$ and Ph_2O give much lower reaction rates. II. *Ibid.*, 1310-13. In the Ulmann-Minaev dehalogenation of 1-chloroanthraquinone in the presence of acetates of alkali metals and a Cu catalyst, the acetate radical suffers decomposition, yielding CO_2 and $\text{HOCH}_2\text{CO}_2\text{H}$. The reduction of chloroanthraquinone proceeds at the expense of the H of K or Na acetates. Since only partial isolation of products has been accomplished, no complete reaction mechanism is proposed at this time. G. M. K.

ARTEM'YEV, N.I.; KRETOV, A.Ye.

Chemism of dehalogenation in the series of 1-halogen derivatives of anthraquinone. Zhur.prikl.khim. 26 no.12:1310-1313 D '53. (MLRA 6:11)

1. Kafedra organicheskoy khimii Dnepropetrovskogo khimiko-tekhnologicheskogo instituta.
(Anthraquinone) (Halogenation)

KRETOV, H. Ye.

Synthesis of arekylamines from chloromethylaryl compounds. A. H. Kretov and E. R. Chertok (Inst. Chem. Technol., Dnepropetrovsk). *Ukrain. Khim. Zvez.* 20, 233-300 (1964) (in Russian). — *p*-Tolylamine, *p*-MeOC₆H₄CH₂NH₂ (I), and *α*-naphthylmethylaniline were prepd. by a new synthesis consisting of (1) chloromethylation, (2) condensation of the CICH₂ derivs. with urea, and (3) decompn. of the condensation products with alkali. Thus, a mixt. of 200 ml. anisole, 200 ml. concd. HCl and 53 g. trioxymethylene agitated 4 hrs. at 8-10°, and the org. layer sepd. and distd. gave 95 g. *p*-CICH₂C₆H₄OMe (II), b_p 76-68°, which when refluxed with urea in H₂O, the mixt. steam distd., the residue sepd., dried, dispersed in boiling H₂O, filtered, the filtrate let stand until crystals appeared (12-16 hrs.), and these filtered off gave 75% *p*-anisylurea. This (10 g.) and 4 g. NaOH mixed, kept 1 hr., and distd. at 240-70° gave 91% I. The other amines were prepd. similarly. CuCl increased catalytically the yield of *N,N'*-bis(*α*-naphthylmethyl)urea. The factors affecting the reactions at the various stages are described in some detail. Gary Gerard

KRETUV, A. Ye.


Subject : USSR/Chemistry AID - P-110
Card : 1/1
Author : Kretov, A. Ye., Dnepropetrovsk
Title : Dicyandiamide and its Reactions
Periodical : Usp. Khim., 23, no. 1, 105-122, 1954
Abstract : The structure and properties of dicyandiamide are
discussed and its reactions thoroughly reviewed.
100 references (6 U.S.S.R.):1862-1951. 1 table.
Institution : None
Submitted : No date

KRETCH, A. E.

✓ Reactivity of chlorine in chlorinated brown coal. M. I. Shenber,
A. E. Kretsch and M. I. Svin (Ukr. Khim. Zh., 1955, 21, 606-
610). Brown coal was treated with Cl_2 (a) in presence of water at
20 or 100°, or (b) in presence of $SnCl_4$ at 100 or 200°, or (c) it was
heated to 325° with HCl in a sealed tube after treatment as in (b).
It was found that only products from (c) were relatively inert to
hydrolytic agents, which indicated that that treatment was necessary
for complete splitting-off of side-chains from the molecules so as
to leave only chlorinated aromatic nuclei. From the results it was
inferred that the degree of aromatization of the brown coal sample
used was 42%.

True

3



1791. ACTIVITY OF CHLORINE IN THE CHLORINATION OF BROWN COAL.
Sienko, M.I., Prato, A.E. and Sovin, M.I. (Ukr. Khim. Zh. (Ukr. Chem. J.),
vol. 21, 535-540; abstr. in Chem. Abstr., 1956, vol. 50, 6771).
Chlorination was carried out in aqueous chlorine solution at 20° and 100°, and

3